arXiv:1404.0491v1 [nlin.SI] 2 Apr 2014

Form factors of the half-infinite XXZ spin chain with a triangular boundary

April 3, 2014

P. Baseilhac [α] and T. Kojima [β]

[α] *Laboratoire de Mathématiques et Physique Théorique CNRS/UMR 7350, Fédération Denis Poisson FR2964, Université de Tours, Parc de Grammont, 37200 Tours, FRANCE*
baseilha@lmpt.univ-tours.fr

[β] *Department of Mathematics and Physics, Faculty of Engineering, Yamagata University, Jonan 4-3-16, Yonezawa 992-8510, JAPAN*
kojima@yz.yamagata-u.ac.jp

Abstract

The half-infinite XXZ spin chain with a triangular boundary is considered in the massive regime. Two integral representations of form factors of local operators are proposed using bosonization. Sufficient conditions such that the expressions for triangular boundary conditions coincide with those for diagonal boundary conditions are identified. The expressions are compared with known results upon specializations. Using the spin-reversal property which relates the Hamiltonian with upper and lower triangular boundary conditions, new identities between multiple integrals of infinite products are extracted.

1 Introduction

Among the family of quantum integrable models, the XXZ spin chain with different types of integrable boundary conditions is the subject of current investigations. For periodic boundary conditions, several results are already known: the spectral problem - eigenvalues and eigenvectors - for the Hamiltonian has been handled either using the Bethe ansatz (BA) approach [1], or using the corner transfer matrix method (CTM) in the thermodynamic limit [2]. In addition, correlation functions and form factors of local operators have been derived using the quantum inverse scattering method (QISM) in [3, 4], or

using the vertex operator approach (VOA) in the thermodynamic limit [1] [5]. It has been checked that expressions of correlation functions obtained by both approaches coincide in the thermodynamic limit. They are given by multiple integrals of meromorphic functions.

For general integrable boundary conditions, less is known. For diagonal [13], triangular [14], related non-diagonal boundaries [15, 16] or q a root of unity [17], the spectral problem of the corresponding finite XXZ open chain has been solved using the BA. Extending the analysis of [4], correlation functions have been also proposed for diagonal boundary conditions [18, 19]. In the thermodynamic limit, the half-infinite XXZ spin chain with general integrable boundary conditions can be alternatively formulated using the VOA, either starting from Sklyanin's framework [13] or from the recently developed Onsager's approach [20, 21] (see also related works [22]). For diagonal and triangular boundary conditions, eigenvalues and eigenvectors are proposed in [23, 21] and integral representations of correlation functions in the massive regime are computed in [23, 24], respectively. In the massless regime and diagonal boundary conditions, note that integral representations are obtained in [25].

Up to now, the calculation of form factors of local operators of the half-infinite XXZ spin chain within the VOA - even in the case of diagonal boundary conditions - has not been addressed. Besides the importance of the form factors in the non-perturbative analysis of this model, there are four main motivations for considering this problem in details. First, an alternative derivation of the QISM proposal for diagonal boundary conditions [18, 19] is highly desirable. Secondly, for *non-diagonal* boundary conditions there are no known results available yet for the form factors, besides the two integral representations of correlation functions recently proposed in [24]. Third, correlation functions and form factors are known to satisfy quantum boundary Knizhnik-Zamolodchikov equations, whose solutions for non-diagonal boundary conditions are recently investigated in [26, 27]. Any new piece of information - for instance integral representations of form factors - is welcome, especially if it is related with[2] the q-Onsager algebra representation theory. Fourth, it is expected that the remarkable identities between the correlation functions given in the form of n-fold multiple integrals of elliptic theta functions and discovered in [24] can be generalized to arbitrary form factors.

In this paper, the half-infinite XXZ spin chain with a triangular boundary is studied in the massive regime. This paper is a continuation of the paper [24]. Integral representations of form factors of local operators are proposed using bosonization. Using the spin-reversal property which relates the Hamiltonian with upper and lower triangular boundary conditions, new identities between multiple integrals of infinite products are extracted. The paper is organized as follows. In Section 2, we briefly recall how the half-infinite XXZ spin chain is formulated within the VOA: eigenvalues, eigenvectors, dual eigenvectors and scalar products between these are given. Integral representations of form factors of local operators for the half-infinite XXZ spin chain with *diagonal* and *triangular* boundary conditions are proposed in Section 3 and Section 4, respectively. In Section 3, upon specializations the expressions are compared

[1] For models associated with higher rank affine Lie algebras, the calculation of correlation functions and form factors using the QISM is currently investigated (see e.g. [6, 7]). In the thermodynamic limit, correlation functions can be derived using the VOA [8, 9, 10, 11, 12].

[2] Type I and type II q-vertex operators entering in the correlation functions and form factors are intertwiners of infinite dimensional (level one) representations of the q-Onsager algebra, a coideal subalgebra of $U_q(\widehat{sl_2})$ [21].

with the known results obtained within the QISM [18] and [23]. Some checks are done in simple cases. Included in Section 3 and Section 4, using the spin-reversal property, relations between multiple integrals of infinite products are extracted. In Section 5 we summarize concluding remarks. In Appendix A, basic definitions of the R-matrix and the K-matrix are recalled. In Appendix B we summarize bosonizations of the vertex operators and normal orderings. In Appendix C we give a direct proof of the spin reversal property for the simplest case. In Appendix D we summarize convenient formulae for the calculations of vacuum expectation values.

2 The half-infinite XXZ chain

In this Section, basic necessary ingredients for the analysis of the form factors of local operators are briefly recalled. We refer the reader to [23, 21, 24] for more details.

2.1 Hamiltonian

In the present paper, we consider the half-infinite XXZ spin chain with Hamiltonian[3]:

$$H_B^{(\pm)} = -\frac{1}{2}\sum_{k=1}^{\infty}(\sigma_{k+1}^x\sigma_k^x + \sigma_{k+1}^y\sigma_k^y + \Delta\sigma_{k+1}^z\sigma_k^z) - \frac{1-q^2}{4q}\frac{1+r}{1-r}\sigma_1^z - \frac{s}{1-r}\sigma_1^{\pm}. \tag{2.2}$$

Here, as usual $\Delta = (q+q^{-1})/2$ denotes the anisotropy parameter. Diagonal boundary conditions correspond to the special case $s = 0$, whereas $s \neq 0$ describes triangular boundary conditions of upper/lower type according to the sign index $(\pm)$ chosen in (2.2). The massive regime corresponds to the range of parameters:

$$-1 < q < 0, \quad -1 \leq r \leq 1, \quad s \in \mathbf{R}. \tag{2.3}$$

Note that under conjugation of $H_B^{(\pm)}$ by the spin-reversal operator $\hat{\nu} = \prod_{j=1}^{\infty}\sigma_j^x$ the sign of the boundary term is reversed. Then, in the discussion below we can restrict our attention to the domain where the boundary term $-\frac{1-q^2}{4q}\frac{1+r}{1-r} \geq 0$, namely $-1 \leq r \leq 1$.

2.2 The transfer matrix in the vertex operator approach

We now turn to the mathematical formulation of the half-infinite XXZ spin chain based on the q-vertex operator approach [23, 21]. In Skylanin's framework the transfer matrix associated with the *finite* XXZ open spin chain is built from a finite number of R-matrices, and two K-matrices [13]. In order to formulate the *infinite* model (2.2), an infinite combination of R-matrices [23] has to be considered. Generally speaking, infinite combinations of the R-matrix are not free from the difficulty of divergences. However the problem can be alternatively tackled using the q-vertex operators introduced by Baxter [30] and

[3]We use the standard Pauli matrices

$$\sigma^x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},\ \sigma^y = \begin{pmatrix} 0 & -\sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix},\ \sigma^z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},\ \sigma^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix},\ \sigma^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \tag{2.1}$$

Jimbo, Miwa, and Nakayashiki [31]. Note that the CTM for $U_q(\widehat{sl_2})$ [32] gives a supporting argument for such mathematical formulation, that we recall now.

Following the strategy of [5, 23], it is possible to formulate the transfer matrix associated with (2.2) using the q-vertex operators of $U_q(\widehat{sl_2})$. Let $\{v_+, v_-\}$ denote the natural basis of $V = \mathbf{C}^2$. Consider the infinite dimensional vector space $\cdots \otimes V_3 \otimes V_2 \otimes V_1$ on which the Hamiltonian (2.2) acts. Let us introduce the subspace $\mathcal{H}^{(i)}$ $(i = 0, 1)$ of the half-infinite spin chain:

$$\mathcal{H}^{(i)} = Span\{\cdots \otimes v_{p(N)} \otimes \cdots \otimes v_{p(2)} \otimes v_{p(1)} |\ p(N) = (-1)^{N+i}\ (N \gg 1)\}, \tag{2.4}$$

where $p : \mathbf{N} \to \{\pm\}$. In the thermodynamic limit, it is conjectured that level one infinite dimensional representations of $U_q(\widehat{sl_2})$ [23, 21] are embedded into $\mathcal{H}^{(i)}$ $(i = 0, 1)$ using the q-vertex operators. Let $V(\Lambda_i)$ $(i = 0, 1)$ denote the level one irreducible highest weight $U_q(\widehat{sl_2})$ representations with fundamental weights Λ_i. Let V_ζ the evaluation representation of the basic representation of $U_q(sl_2)$. In the VOA we identify the space $\mathcal{H}^{(i)}$ with the space $V(\Lambda_i)$:

$$\mathcal{H}^{(i)} = V(\Lambda_i). \tag{2.5}$$

Introduce the type-I vertex operators $\Phi_\epsilon^{(1-i,i)}(\zeta)$ as the intertwiner of $U_q(\widehat{sl_2})$ [5, 23].

$$\Phi^{(1-i,i)}(\zeta) : V(\Lambda_i) \to V(\Lambda_{1-i}) \otimes V_\zeta, \quad \Phi^{(1-i,i)}(\zeta) = \sum_\epsilon \Phi_\epsilon^{(1-i,i)}(\zeta) \otimes v_\epsilon. \tag{2.6}$$

They satisfy the following commutation relation, the duality, and the invertibility properties:

$$\Phi_{\epsilon_2}^{(i,1-i)}(\zeta_2)\Phi_{\epsilon_1}^{(1-i,i)}(\zeta_1) = \sum_{\epsilon_1',\epsilon_2'=\pm} R(\zeta_1/\zeta_2)_{\epsilon_1\epsilon_2}^{\epsilon_1'\epsilon_2'} \Phi_{\epsilon_1'}^{(i,1-i)}(\zeta_1)\Phi_{\epsilon_2'}^{(1-i,i)}(\zeta_2), \tag{2.7}$$

$$\Phi_\epsilon^{*(1-i,i)}(\zeta) = \Phi_{-\epsilon}^{(1-i,i)}(-q^{-1}\zeta), \tag{2.8}$$

$$g\sum_{\epsilon=\pm} \Phi_\epsilon^{*(i,1-i)}(\zeta)\Phi_\epsilon^{(1-i,i)}(\zeta) = id, \quad g\Phi_{\epsilon_1}^{(i,1-i)}(\zeta)\Phi_{\epsilon_2}^{*(1-i,i)}(\zeta) = \delta_{\epsilon_1\epsilon_2} id, \tag{2.9}$$

where the $R-$matrix $R(\zeta)$ is given in Appendix A, and

$$g = \frac{(q^2;q^4)_\infty}{(q^4;q^4)_\infty}, \qquad (z;p)_\infty = \prod_{n=0}^{\infty}(1 - p^n z). \tag{2.10}$$

As the generating function of the Hamiltonian $H_B^{(\pm)}$ (2.2) one introduces the "renormalized" transfer matrix $T_B^{(\pm,i)}(\zeta; r, s)$ using the q-vertex operators [23, 21]:

$$T_B^{(\pm,i)}(\zeta; r, s) = g \sum_{\epsilon_1,\epsilon_2=\pm} \Phi_{\epsilon_1}^{*(i,1-i)}(\zeta^{-1}) K^{(\pm)}(\zeta; r, s)_{\epsilon_1}^{\epsilon_2} \Phi_{\epsilon_2}^{(1-i,i)}(\zeta) \tag{2.11}$$

where the K-matrices $K^{(\pm)}(\zeta)$ are reported in Appendix A. In the VOA we relate the Hamiltonian $H_B^{(\pm)}$ to the "renormalized" transfer matrix $T_B^{(\pm,i)}(\zeta; r, s)$ by

$$\frac{d}{d\zeta} T_B^{(\pm,i)}(\zeta; r, s)\Big|_{\zeta=1} = \frac{4q}{1-q^2} H_B^{(\pm)} + \text{const.} \tag{2.12}$$

From the boundary Yang-Baxter equation satisfied by $K^{(\pm)}(\zeta)$ and the properties of the q-vertex operators (A.8), (A.9), (2.7), (2.8), (2.9), the following properties of the "renormalized" transfer matrix hold:

$$[T_B^{(\pm,i)}(\zeta_1;r,s), T_B^{(\pm,i)}(\zeta_2;r,s)] = 0 \quad \text{for any } \zeta_1,\zeta_2, \tag{2.13}$$

$$T_B^{(\pm,i)}(1;r,s) = id, \quad T_B^{(\pm,i)}(\zeta;r,s) T_B^{(\pm,i)}(\zeta^{-1};r,s) = id, \tag{2.14}$$

$$T_B^{(\pm,i)}(-q^{-1}\zeta^{-1};r,s) = T_B^{(\pm,i)}(\zeta;r,s). \tag{2.15}$$

Note that the CTM argument for $U_q(\widehat{sl_2})$ [32], as well as the alternative support within Onsager's framework [21], justify the introduction of the transfer matrix (2.11). Besides the fact that the q-vertex operators $\Phi_\epsilon^{(1-i,i)}(\zeta)$, $\Phi_\epsilon^{*(1-i,i)}(\zeta)$ associated with $U_q(\widehat{sl_2})$ are well-defined objects - free from the difficulty of divergences - they are the unique solution of the intertwining relations defining the q-vertex operators of the q-Onsager algebra, which characterizes the hidden non-Abelian symmetry of (2.2) (see [21] for details).

2.3 Eigenvectors, duals and basic scalar products

For the model (2.2) with diagonal ($s = 0$) or triangular ($s \neq 0$) boundary conditions, finding the eigenvalues, eigenvectors and dual eigenvectors of the Hamiltonian leads to consider the spectral problem of the transfer matrix (2.11). In the VOA, two fundamental solutions - the so-called vacuum eigenvectors $|\pm;i\rangle_B$ ($i = 0, 1$) - can be constructed using bosonization. Dual vacuum eigenvectors ${}_B\langle i;\pm|$ are constructed similarly. For general values s, by definition they satisfy

$$T_B^{(\pm,i)}(\zeta;r,s)|\pm;i\rangle_B = \Lambda^{(i)}(\zeta;r)|\pm;i\rangle_B, \tag{2.16}$$

$${}_B\langle i;\pm|T_B^{(\pm,i)}(\zeta;r,s) = {}_B\langle i;\pm|\Lambda^{(i)}(\zeta;r)\ . \tag{2.17}$$

Proposed in [23] for $s = 0$ and [24] for $s \neq 0$, the structure of the eigenvectors and duals is as follows. Let e_j, f_j, h_j, $j = 0, 1$, be the Chevalley generators of $U_q(\widehat{sl_2})$. The vacuum eigenvectors read[4]:

$$|+;0\rangle_B = \exp_{q^{-1}}(sf_0)\,|0\rangle_B, \qquad |+;1\rangle_B = \exp_q\left(\frac{s}{rq}e_1 q^{-h_1}\right)|1\rangle_B, \tag{2.18}$$

$$|-;0\rangle_B = \exp_q\left(-\frac{s}{q}e_0 q^{-h_0}\right)|0\rangle_B, \qquad |-;1\rangle_B = \exp_{q^{-1}}\left(-\frac{s}{r}f_1\right)|1\rangle_B, \tag{2.19}$$

where $|i\rangle_B$ are the vacuum eigenvectors of the diagonal boundary $s = 0$ [23] (see definitions below). Dual vacuum eigenvectors take the form:

$${}_B\langle 0;+| = {}_B\langle 0|\exp_q(-sf_0), \qquad {}_B\langle 1;+| = {}_B\langle 1|\exp_{q^{-1}}\left(-\frac{s}{rq}e_1 q^{-h_1}\right), \tag{2.20}$$

$${}_B\langle 0;-| = {}_B\langle 0|\exp_{q^{-1}}\left(\frac{s}{q}e_0 q^{-h_0}\right), \qquad {}_B\langle 1;-| = {}_B\langle 1|\exp_q\left(\frac{s}{r}f_1\right), \tag{2.21}$$

where ${}_B\langle i|$ are the dual vacuum eigenvectors of the diagonal boundary $s = 0$ [23]. For $s = 0$, realizations of vacuum vectors and dual ones in terms of q-bosons have been obtained in [23]. See the definition of

[4]Here we have introduced the q-exponential function $\exp_q(x)$ defined by $\exp_q(x) = \sum_{n=0}^{\infty} \frac{q^{\frac{n(n-1)}{2}}}{[n]_q!}x^n$.

the q-bosons a_n ($n \in \mathbf{N}_{\neq 0}$) in Appendix B. Let us denote $|i\rangle$ (resp. $\langle i|$) the highest (resp. dual highest) weight vector by:

$$|i\rangle = 1 \otimes e^{\Lambda_i}, \qquad \langle i| = 1 \otimes e^{-\Lambda_i}, \tag{2.22}$$

respectively. Explicitly, for the diagonal case $s = 0$ the vacuum eigenvectors and duals are given by:

$$|i\rangle_B = \exp(F_i)\,|i\rangle, \qquad F_i = \frac{1}{2}\sum_{n=1}^{\infty} \frac{n\alpha_n}{[2n]_q[n]_q} a_{-n}^2 + \sum_{n=1}^{\infty} \beta_n^{(i)} a_{-n}, \tag{2.23}$$

$${}_B\langle i| = \langle i|\exp(G_i), \qquad G_i = \frac{1}{2}\sum_{n=1}^{\infty} \frac{n\gamma_n}{[2n]_q[n]_q} a_n^2 + \sum_{n=1}^{\infty} \delta_n^{(i)} a_n. \tag{2.24}$$

Here we have set

$$\alpha_n = -q^{6n}, \quad \gamma_n = -q^{-2n}, \tag{2.25}$$

and

$$\beta_n^{(i)} = -\theta_n \frac{q^{5n/2}(1-q^n)}{[2n]_q} + \begin{cases} -\frac{q^{7n/2}r^n}{[2n]_q} & (i=0), \\ +\frac{q^{3n/2}r^{-n}}{[2n]_q} & (i=1), \end{cases} \tag{2.26}$$

$$\delta_n^{(i)} = \theta_n \frac{q^{-3n/2}(1-q^n)}{[2n]_q} + \begin{cases} -\frac{q^{-5n/2}r^n}{[2n]_q} & (i=0), \\ +\frac{q^{-n/2}r^{-n}}{[2n]_q} & (i=1), \end{cases} \tag{2.27}$$

where

$$[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \qquad \theta_n = \begin{cases} 1 & \text{for } n \text{ even}, \\ 0 & \text{for } n \text{ odd}. \end{cases} \tag{2.28}$$

In addition, the two fundamental eigenvalues of the transfer matrix are given by[5]

$$\Lambda^{(i)}(\zeta; r) = \begin{cases} 1 & (i=0) \\ \frac{1}{\zeta^2}\frac{\Theta_{q^4}(r\zeta^2)\Theta_{q^4}(q^2 r\zeta^{-2})}{\Theta_{q^4}(r\zeta^{-2})\Theta_{q^4}(q^2 r\zeta^2)}, & (i=1) \end{cases}. \tag{2.29}$$

Starting from the two vacuum eigenvectors, it is possible to create the excited states by an application of type-II vertex operators. Recall that type-II vertex operators $\Psi_\epsilon^{*(1-i,i)}(\xi)$ are also intertwiners of $U_q(\widehat{sl_2})$. Together with the type-I vertex operators $\Phi_\epsilon^{(1-i,i)}(\zeta)$, the type-II vertex operators $\Psi_\mu^{(i,1-i)}(\xi)$ satisfy the following commutation relation:

$$\Phi_\epsilon^{(i,1-i)}(\zeta)\Psi_\mu^{*(1-i,i)}(\xi) = \tau(\zeta/\xi)\Psi_\mu^{*(i,1-i)}(\xi)\Phi_\epsilon^{(1-i,i)}(\zeta), \tag{2.30}$$

with

$$\tau(\zeta) = \zeta^{-1}\frac{\Theta_{q^4}(q\zeta^2)}{\Theta_{q^4}(q\zeta^{-2})}. \tag{2.31}$$

This commutation relation implies that

$$\Psi_{\mu_1}^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_{\mu_{N-1}}^{*(i+2,i+1)}(\xi_{N-1})\Psi_{\mu_N}^{*(i+1,i)}(\xi_N)|\pm; i\rangle_B \tag{2.32}$$

[5]The elliptic theta function defined by $\Theta_p(z) = (p;p)_\infty(z;p)_\infty(p/z;p)_\infty$ is introduced.

is an eigenvector of $T_B^{(\pm,i)}(\zeta;r,s)$ with eigenvalue :

$$\Lambda^{(i)}(\zeta;r)\prod_{j=1}^{N}\tau(\zeta/\xi_j)\tau(\zeta\xi_j). \tag{2.33}$$

Here the number $i+N$ in the suffix $(i+N,i+N-1)$ of the q-vertex operator $\Psi_{\mu_1}^{*(i+N,i+N-1)}(\xi_1)$ should be understood modulo 2. Note that type-II vertex operators are intertwiners of the q-Onsager algebra, which also justify their use in solving (2.2) [21].

Finally, from the relation for the q-exponential function $\exp_q(x)\exp_{q^{-1}}(-x)=1$ the basic scalar products between the vacuum eigenvectors and their duals can be computed. Observe:

$$ {}_B\langle i;\pm|\pm;i\rangle_B = {}_B\langle i|i\rangle_B. \tag{2.34}$$

Hence, according to the formulae for the norms of ${}_B\langle i|i\rangle_B$ given in [23] it follows:

$$ {}_B\langle i;\pm|\pm;i\rangle_B = \begin{cases} \frac{(q^4r^2;q^8)_\infty}{(q^6;q^8)_\infty(q^2r^2;q^8)_\infty} & (i=0), \\ \frac{(q^4/r^2;q^8)_\infty}{(q^6;q^8)_\infty(q^2/r^2;q^8)_\infty} & (i=1). \end{cases} \tag{2.35}$$

2.4 Spin-reversal property

In the thermodynamic limit, the analogue in the VOA of the spin-reversal operator $\hat{\nu}:\mathcal{H}^{(i)}\longrightarrow\mathcal{H}^{(1-i)}$ $(i=0,1)$ is played by the vector-space isomorphism $\nu: V(\Lambda_i)\longrightarrow V(\Lambda_{1-i})$, associated with the Dynkin diagram symmetry [5]. Recall that the action of ν on the Chevalley generators of $U_q(\widehat{sl_2})$ is given by:

$$\nu e_j\nu = e_{1-j},\quad \nu f_j\nu = f_{1-j},\quad \nu q^{h_j}\nu = q^{h_{1-j}}. \tag{2.36}$$

Moreover, one has [5]:

$$\nu\ \Phi_\epsilon^{(i,1-i)}(\zeta)\ \nu = \Phi_{-\epsilon}^{(1-i,i)}(\zeta),\qquad \nu\ \Psi_\mu^{*(i,1-i)}(\zeta)\ \nu = \Psi_{-\mu}^{*(1-i,i)}(\zeta). \tag{2.37}$$

Under the action of ν, it is possible to show that the transfer matrix (2.11) satisfies (see [23, 24] for details):

$$\nu T_B^{(\pm,1)}(\zeta;r,s)\nu = \Lambda^{(1)}(\zeta;r)T_B^{(\mp,0)}(\zeta;1/r,-s/r). \tag{2.38}$$

As a consequence, according to the structure of (2.18)-(2.21) and the results for the diagonal case $s=0$ [23], the vacuum eigenvectors and their duals should be related by[6]:

$$\nu\left(|\pm;i\rangle_B\right)|_{r\to1/r,s\to-s/r} = |\mp;1-i\rangle_B, \tag{2.39}$$

$$\nu\left({}_B\langle i;\pm|\right)|_{r\to1/r,s\to-s/r} = {}_B\langle 1-i;\mp|. \tag{2.40}$$

Above relations will be used in further Sections, for instance to derive two integrals representations for the form factors, as well as linear the relations between multiple integrals.

[6] A rigorous proof of these relations - besides the known consistency checks - in the bosonization framework remains an open problem.

3 Form factors for a diagonal boundary

In this Section, we study the form factors of local operators for a *diagonal* boundary. Integral representations of form factors of local operators are proposed using bosonization.

3.1 Definition

In this Section we study the vacuum expectation values of the q-vertex operators with natural numbers M, N with $M = N \ (mod.2)$:

$$\begin{aligned} & F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r) \\ = & \frac{{}_B\langle i|\Phi^{(i,i+1)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(i+M-1,i+M)}_{\epsilon_M}(\zeta_M)\Psi^{*(i+N,i+N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(i+1,i)}_{\mu_N}(\xi_N)|i\rangle_B}{{}_B\langle i|i\rangle_B}. \end{aligned} \tag{3.1}$$

Here the number i in the suffix $(i, i+1)$ of q-vertex operators $\Phi^{(i,i+1)}_{\epsilon}(\zeta)$ should be understood modulo 2.

Let L be a linear operator on the n-fold tensor products of the two-dimensional space $V_n \otimes \cdots \otimes V_2 \otimes V_1$. The corresponding local operator $\mathcal{L}$ acting on $V(\Lambda_i)$ can be defined in terms of the type-I vertex operators in exactly the same way as in the bulk theory [5]. Explicitly, if $(E_{\epsilon\epsilon'})_n$ is the matrix at the n-th site $E_{\epsilon\epsilon'} \otimes \underbrace{id \otimes \cdots \otimes id}_{n-1}$, the corresponding local operator $(\mathcal{E}_{\epsilon\epsilon'})_n$ is given by $(\mathcal{E}_{\epsilon\epsilon'})_n = E_{\epsilon\epsilon'}(1, 1, \cdots, 1)$, where we have set

$$\begin{aligned} E_{\epsilon\epsilon'}(\zeta_1,\zeta_2,\cdots,\zeta_n) &= g^n \sum_{\epsilon_1,\cdots,\epsilon_{n-1}=\pm} \Phi^{*(i,i+1)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{*(i+n-2,i+n-1)}_{\epsilon_{n-1}}(\zeta_{n-1})\Phi^{*(i+n-1,i+n)}_{\epsilon}(\zeta_n) \\ &\times \Phi^{(i+n,i+n-1)}_{\epsilon'}(\zeta_n)\Phi^{(i+n-2,i+n-1)}_{\epsilon_{n-1}}(\zeta_{n-1})\cdots\Phi^{(i+1,i)}_{\epsilon_1}(\zeta_1). \end{aligned} \tag{3.2}$$

From the inversion property of the q-vertex operators (2.9), the form factors of local operators $(\mathcal{E}_{\epsilon_m\epsilon'_m})_m(\mathcal{E}_{\epsilon_{m-1}\epsilon'_{m-1}})_{m-1}\cdots(\mathcal{E}_{\epsilon_1\epsilon'_1})_1$ with $M = 2m$ and N : even are given by

$$\begin{aligned} & {}_B\langle i|(\mathcal{E}_{\epsilon_m\epsilon'_m})_m(\mathcal{E}_{\epsilon_{m-1}\epsilon'_{m-1}})_{m-1}\cdots(\mathcal{E}_{\epsilon_1\epsilon'_1})_1\Psi^{*(i+N,i+N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(i+1,i)}_{\mu_N}(\xi_N)|i\rangle_B \\ = & g^m \times {}_B\langle i|i\rangle_B \times F^{(i,2m,N)}_{-\epsilon_1,\cdots,-\epsilon_m,\epsilon'_m,\cdots,\epsilon'_1,\mu_1,\cdots,\mu_N}(\overbrace{-q^{-1},\cdots,-q^{-1}}^{m},\overbrace{1,\cdots,1}^{m},\xi_1,\cdots,\xi_N;r). \end{aligned} \tag{3.3}$$

In what follows we call both (3.1) and (3.3) the form factors of local operators. In the next Section we calculate the form factors by using bosonizations.

3.2 Integral representation

In this Section we focus our attention on the half-infinite XXZ spin chain with a diagonal boundary. We calculate the form factors with $M = N \ (mod.2)$:

$$F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r). \tag{3.4}$$

Here we summarize detailed calculations. In this Section we set

$$A = \{1 \le a \le M | \epsilon_a = +\}, \quad B = \{1 \le b \le N | \mu_b = +\}. \tag{3.5}$$

Let $|A|$ and $|B|$ denote number of elements of the sets A and B, respectively. Note that the following infinite product relation :

$$\exp\left(-\sum_{n=1}^{\infty}\frac{1}{n}\frac{z^n}{(1-p_1^n)(1-p_2^n)\cdots(1-p_N^n)}\right)=(z;p_1,p_2,\cdots,p_N)_\infty \tag{3.6}$$

will be used, where we denote

$$(z;p_1,p_2,\cdots,p_N)_\infty=\prod_{n_1,n_2,\cdots,n_N=0}^{\infty}(1-p_1^{n_1}p_2^{n_2}\cdots p_N^{n_N}z). \tag{3.7}$$

We introduce the double-infinite products :

$$\{z\}_\infty=(z;q^4,q^4)_\infty,\qquad [z]_\infty=(z;q^8,q^8)_\infty. \tag{3.8}$$

Extending the analysis of [24], the normal ordering of products of type-I and type-II vertex operators are given by

$$\begin{aligned}
&\Phi_{\epsilon_1}^{(i,i+1)}(\zeta_1)\cdots\Phi_{\epsilon_M}^{(i+M-1,i+M)}(\zeta_M)\cdot\Psi_{\mu_1}^{*(i+N-1,i+N)}(\xi_1)\cdots\Psi_{\mu_N}^{*(i+1,i)}(\xi_N)\\
=\;& q^{|B|}(-q^3)^{\frac{1}{4}M(M-1)+\frac{1}{4}N(N-1)-\frac{1}{2}MN-\sum_{a\in A}a-\sum_{b\in B}b}(1-q^2)^{|A|+|B|}\prod_{j=1}^{M}\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j}\prod_{j=1}^{N}\xi_j^{\frac{1+\mu_j}{2}+N-j}\\
\times\;&\prod_{1\le j<k\le M}\frac{(q^2\zeta_k^2/\zeta_j^2;q^4)_\infty}{(q^4\zeta_k^2/\zeta_j^2;q^4)_\infty}\prod_{1\le j<k\le N}\frac{(\xi_k^2/\xi_j^2;q^4)_\infty}{(q^2\xi_k^2/\xi_j^2;q^4)_\infty}\prod_{j=1}^{M}\prod_{k=1}^{N}\frac{(q^3\xi_k^2/\zeta_j^2;q^4)_\infty}{(q\xi_k^2/\zeta_j^2;q^4)_\infty}\\
\times\;&\oint\cdots\oint_C\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a\prod_{b\in B}\frac{du_b}{2\pi\sqrt{-1}}\frac{\prod_{\substack{a,b\in A\\a<b}}(w_a-w_b)(w_a-q^2w_b)\prod_{\substack{a,b\in B\\a<b}}(u_a-u_b)(u_a-q^{-2}u_b)}{\prod_{a\in A}\prod_{b\in B}(w_a-qu_b)(w_a-q^{-1}u_b)}\\
\times\;&\frac{\prod_{a\in A}\prod_{j=1}^{N}(w_a-q^3\xi_j^2)\prod_{b\in B}\prod_{j=1}^{M}(u_b-q^3\zeta_j^2)}{\prod_{a\in A}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}\prod_{b\in B}\left\{\prod_{1\le j\le b}(\xi_j^2-q^{-4}u_b)\prod_{b\le j\le N}(u_b-q^2\xi_j^2)\right\}}\\
\times\;&:\Phi_-^{(i,i+1)}(\zeta_1)\cdots\Phi_-^{(i+M-1,i+M)}(\zeta_M)\Psi_-^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_-^{*(i+1,i)}(\xi_N)\prod_{a\in A}X^-(w_a)\prod_{b\in B}X^+(u_b):.
\end{aligned} \tag{3.9}$$

Here the integral contour C is simple closed curve such that the w_a $(a\in A)$ encircles $q^4\zeta_j^2$ $(a\le j\le M)$, $qu_b, q^{-1}u_b$ $(b\in B)$ but not $q^2\zeta_j^2$ $(1\le j\le a)$, and that the u_b $(b\in B)$ encircles $q^2\xi_j^2$ $(b\le j\le N)$ but not $q^4\xi_j^2$ $(1\le j\le b)$, $qw_a, q^{-1}w_a$ $(a\in A)$. Normal orderings of the basic operators are summarized in Appendix B. The zero-mode e^α part of the operator :

$$:\Phi_-^{(i,i+1)}(\zeta_1)\cdots\Phi_-^{(i+M-1,i+M)}(\zeta_M)\Psi_-^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_-^{*(i+1,i)}(\xi_N)\prod_{a\in A}X^-(w_a)\prod_{b\in B}X^+(u_b):$$

is given by $e^{\alpha((M-N)+2(|B|-|A|))}$. Then, the condition for which the vacuum expectation value is non-vanishing reads :

$$\begin{aligned}
&{}_B\langle i|:\prod_{j=1}^{M}\Phi_-^{(i+j-1,i+j)}(\zeta_j)\prod_{j=1}^{N}\Psi_-^{*(i+j-1,i+j)}(\xi_j)\prod_{a\in A}X^-(w_a)\prod_{b\in B}X^+(u_b):|i\rangle_B\neq 0,\\
\Longleftrightarrow\;&{}_B\langle i|e^{\alpha((M-N)+2(|B|-|A|))}|i\rangle_B\neq 0\Longleftrightarrow(M-N)=2(|A|-|B|).
\end{aligned} \tag{3.10}$$

Provided the condition $(M-N)=2(|A|-|B|)$ is satisfied, the vacuum expectation values of the q-vertex operators are given by

$$
\begin{aligned}
& {}_B\langle i|\Phi_{\epsilon_1}^{(i,i+1)}(\zeta_1)\cdots\Phi_{\epsilon_M}^{(i+M-1,i+M)}(\zeta_M)\Psi_{\mu_1}^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_{\mu_{2N}}^{*(i+1,i)}(\xi_N)|i\rangle_B \\
= \;& q^{|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}(1-q^2)^{|A|+|B|}\prod_{j=1}^{M}\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i}\prod_{j=1}^{N}\xi_j^{\frac{1+\mu_j}{2}+N-j+1-i} \\
\times \;& \prod_{1\le j<k\le M}\frac{(q^2\zeta_k^2/\zeta_j^2;q^4)_\infty}{(q^4\zeta_k^2/\zeta_j^2;q^4)_\infty}\prod_{1\le j<k\le N}\frac{(\xi_k^2/\xi_j^2;q^4)_\infty}{(q^2\xi_k^2/\xi_j^2;q^4)_\infty}\prod_{j=1}^{M}\prod_{k=1}^{N}\frac{(q^3\xi_k^2/\zeta_j^2;q^4)_\infty}{(q\xi_k^2/\zeta_j^2;q^4)_\infty} \\
\times \;& \oint\cdots\oint_C\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a^{1-i}\prod_{a\in B}\frac{du_a}{2\pi\sqrt{-1}}u_a^{i}\frac{\prod_{\substack{a,b\in A\\ a<b}}(w_a-w_b)(w_a-q^2w_b)\prod_{\substack{a,b\in B\\ a<b}}(u_a-u_b)(u_a-q^{-2}u_b)}{\prod_{a\in A}\prod_{b\in B}(w_a-qu_b)(w_a-q^{-1}u_b)} \\
\times \;& \frac{\prod_{a\in A}\prod_{j=1}^{N}(w_a-q^3\xi_j^2)\prod_{a\in B}\prod_{j=1}^{M}(u_a-q^3\zeta_j^2)}{\prod_{a\in A}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}\prod_{a\in B}\left\{\prod_{1\le j\le a}(\xi_j^2-q^{-4}u_a)\prod_{a\le j\le N}(u_a-q^2\xi_j^2)\right\}} \\
\times \;& {}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^-(w_a)+\sum_{a\in B}R^+(u_a)} \\
\times \;& e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^-(w_a)+\sum_{a\in B}S^+(v_a)}|i\rangle_B. \qquad (3.11)
\end{aligned}
$$

Next we calculate the following vacuum expectation value explicitly.

$$
\begin{aligned}
& {}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^-(w_a)+\sum_{a\in B}R^+(u_a)} \qquad (3.12) \\
\times \;& e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^-(w_a)+\sum_{a\in B}S^+(v_a)}|i\rangle_B=\langle i|e^{G^{(i)}}e^{\sum_{n=1}^{\infty}a_{-n}X_n}e^{-\sum_{n=1}^{\infty}a_nY_n}e^{F^{(i)}}|i\rangle.
\end{aligned}
$$

Here we have set

$$
X_n = \frac{q^{7n/2}}{[2n]}\sum_{j=1}^{M}\zeta_j^{2n}-\frac{q^{5n/2}}{[2n]}\sum_{j=1}^{N}\xi_j^{2n}-\frac{q^{n/2}}{[n]}\sum_{a\in A}w_a^n+\frac{q^{-n/2}}{[n]}\sum_{a\in B}u_a^n, \qquad (3.13)
$$

$$
Y_n = \frac{q^{-5n/2}}{[2n]}\sum_{j=1}^{M}\zeta_j^{-2n}-\frac{q^{-7n/2}}{[2n]}\sum_{j=1}^{N}\xi_j^{-2n}-\frac{q^{n/2}}{[n]}\sum_{a\in A}w_a^{-n}+\frac{q^{-n/2}}{[n]}\sum_{a\in B}u_a^{-n}. \qquad (3.14)
$$

We use the following formula given in [23].

$$
\begin{aligned}
& \frac{{}_B\langle i|\exp\left(\sum_{n=1}^{\infty}a_{-n}X_n\right)\exp\left(-\sum_{n=1}^{\infty}a_nY_n\right)|i\rangle_B}{{}_B\langle +;i|i;+\rangle_B} \qquad (3.15) \\
= \;& \exp\left(\sum_{n=1}^{\infty}\frac{[2n][n]}{n}\frac{1}{1-\alpha_n\gamma_n}\left\{\frac{1}{2}\gamma_nX_n^2-\alpha_n\gamma_nX_nY_n+\frac{1}{2}\alpha_nY_n^2+(\delta_n^{(i)}+\gamma_n\beta_n^{(i)})X_n-(\beta_n^{(i)}+\alpha_n\delta_n^{(i)})Y_n\right\}\right).
\end{aligned}
$$

Here we have used α_n,γ_n, $\beta_n^{(i)}$, and $\delta_n^{(i)}$ in (2.25), (2.26), and (2.27), respectively. The formulae summarized in Appendix D are useful for calculations.

• The following is the main result of this Section. For $(|A|-|B|)=\frac{1}{2}(M-N)$, we have the following integral representation of the form factors of local operators :

$$
\begin{aligned}
& F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots.\zeta_M,\xi_1,\cdots,\xi_N;r) \\
= \;& q^{|B|}(1-q^2)^{|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}\left(\frac{\{q^6\}_\infty}{\{q^8\}_\infty}\right)^M\left(\frac{\{q^4\}_\infty}{\{q^6\}_\infty}\right)^N\{(q^2;q^2)_\infty\}^{|A|+|B|}
\end{aligned}
$$

$$
\begin{aligned}
&\times \prod_{1\le j<k\le M} \frac{\{q^6\zeta_j^2\zeta_k^2\}_\infty\{q^2/\zeta_j^2\zeta_k^2\}_\infty\{q^6\zeta_j^2/\zeta_k^2\}_\infty\{q^2\zeta_k^2/\zeta_j^2\}_\infty}{\{q^8\zeta_j^2\zeta_k^2\}_\infty\{q^4/\zeta_j^2\zeta_k^2\}_\infty\{q^8\zeta_j^2/\zeta_k^2\}_\infty\{q^4\zeta_k^2/\zeta_j^2\}_\infty} \\
&\times \prod_{1\le j<k\le N} \frac{\{q^4\xi_j^2\xi_k^2\}_\infty\{1/\xi_j^2\xi_k^2\}_\infty\{q^4\xi_j^2/\xi_k^2\}_\infty\{\xi_k^2/\xi_j^2\}_\infty}{\{q^6\xi_j^2\xi_k^2\}_\infty\{q^2/\xi_j^2\xi_k^2\}_\infty\{q^6\xi_j^2/\xi_k^2\}_\infty\{q^2\xi_k^2/\xi_j^2\}_\infty} \\
&\times \prod_{j=1}^{M}\prod_{k=1}^{N} \frac{\{q^7\zeta_j^2\xi_k^2\}_\infty\{q^3/\zeta_j^2\xi_k^2\}_\infty\{q^7\zeta_j^2/\xi_k^2\}_\infty\{q^3\xi_k^2/\zeta_j^2\}_\infty}{\{q^5\zeta_j^2\xi_k^2\}_\infty\{q/\zeta_j^2\xi_k^2\}_\infty\{q^5\zeta_j^2/\xi_k^2\}_\infty\{q\xi_k^2/\zeta_j^2\}_\infty} \\
&\times \prod_{j=1}^{M} \frac{[q^{10}\zeta_j^4]_\infty[q^{14}\zeta_j^4]_\infty[q^{10}/\zeta_j^4]_\infty[q^6/\zeta_j^4]_\infty}{[q^{12}\zeta_j^4]_\infty[q^{16}\zeta_j^4]_\infty[q^{12}/\zeta_j^4]_\infty[q^8/\zeta_j^4]_\infty} \prod_{j=1}^{N} \frac{[q^4\xi_j^4]_\infty[q^8\xi_j^4]_\infty[1/\xi_j^4]_\infty[q^4/\xi_j^4]_\infty}{[q^6\xi_j^4]_\infty[q^{10}\xi_j^4]_\infty[q^2/\xi_j^4]_\infty[q^6/\xi_j^4]_\infty} \\
&\times \prod_{j=1}^{M} \zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i} \prod_{j=1}^{N} \xi_j^{\frac{\mu_j+1}{2}+N-j+1-i} \oint\cdots\oint_{C^{(i)}} \prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}} w_a^{1-i} \prod_{b\in B}\frac{du_b}{2\pi\sqrt{-1}} u_b^i \\
&\times \frac{\prod_{a\in A}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty \prod_{b\in B}(u_b^2/q^2;q^4)_\infty(q^6/u_b^2;q^4)_\infty}{\prod_{a\in A}\prod_{b\in B} w_a^2(w_au_b/q^3;q^2)_\infty(q^3w_a/u_b;q^2)_\infty(u_b/qw_a;q^2)_\infty(q^5/w_au_b;q^2)_\infty} \\
&\times \frac{\prod_{\substack{a,b\in A\\ a<b}} w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a\in A}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}} \\
&\times \frac{\prod_{\substack{a,b\in B\\ a<b}} u_a^2(u_au_b/q^4;q^2)_\infty(q^2u_a/u_b;q^2)_\infty(u_b/q^2u_a;q^2)_\infty(q^4/u_au_b;q^2)_\infty}{\prod_{b\in B}\left\{\prod_{1\le j\le b}(\xi_j^2-q^{-4}u_b)\prod_{b\le j\le M}(u_b-q^2\xi_j^2)\right\}} \\
&\times \frac{\prod_{j=1}^{M}\prod_{b\in B} u_b(q\zeta_j^2u_b;q^4)_\infty(q^3\zeta_j^2/u_b;q^4)_\infty(qu_b/\zeta_j^2;q^4)_\infty(q^3/\zeta_j^2u_b;q^4)_\infty}{\prod_{j=1}^{M}\prod_{a\in A}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta_j^2;q^4)_\infty(q^4/\zeta_j^2w_a;q^4)_\infty} \\
&\times \frac{\prod_{j=1}^{N}\prod_{a\in A} w_a(q\xi_j^2w_a;q^4)_\infty(q^3\xi_j^2/w_a;q^4)_\infty(qw_a/\xi_j^2;q^4)_\infty(q^3/\xi_j^2w_a;q^4)_\infty}{\prod_{j=1}^{N}\prod_{b\in B}(\xi_j^2u_b;q^4)_\infty(q^6\xi_j^2/u_b;q^4)_\infty(u_b/\xi_j^2;q^4)_\infty(q^2/\xi_j^2u_b;q^4)_\infty} \\
&\times \begin{cases} \prod_{j=1}^{M}\frac{(q^2r\zeta_j^2;q^4)_\infty}{(q^4r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q^3r\xi_j^2;q^4)_\infty}{(qr\xi_j^2;q^4)_\infty}\frac{\prod_{b\in B}(1-ru_b/q^3)}{\prod_{a\in A}(1-rw_a/q^2)} & (i=0), \\ \prod_{j=1}^{M}\frac{(1/r\zeta_j^2;q^4)_\infty}{(q^2/r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q/r\xi_j^2;q^4)_\infty}{(1/qr\xi_j^2;q^4)_\infty}\frac{\prod_{b\in B}(1-q/ru_b)}{\prod_{a\in A}(1-q^2/rw_a)} & (i=1). \end{cases}
\end{aligned}
\tag{3.16}
$$

Here the integration contour $C^{(0)}$ is a closed curve that satisfies the following conditions for $s=0,1,2,\cdots$. The w_a $(a\in A)$ encircles $q^{8+4s}\zeta_j^2$ $(1\le j<a)$, $q^{4+4s}\zeta_j^2$ $(a\le j\le M)$, q^{4+4s}/ζ_j^2 $(1\le j\le M)$ $q^{-1+2s}u_b$, q^{5+2s}/u_b $(b\in B)$, but not $q^{2-4s}\zeta_j^2$ $(1\le j\le a)$, $q^{-2-4s}\zeta_j^2$ $(a<j\le M)$, q^{-2-4s}/ζ_j^2 $(1\le j\le M)$, $q^{-3-2s}u_b$, q^{3-2s}/u_b $(b\in B)$, and q^2/r. The u_b $(b\in B)$ encircles $q^{6+4s}\xi_j^2$ $(1\le j<b)$, $q^{2+4s}\xi_j^2$ $(b\le j\le N)$, q^{2+4s}/ξ_j^2 $(1\le j\le N)$, $q^{3+2s}w_a$, q^{5+2s}/w_a $(a\in B)$, but not $q^{4-4s}\xi_j^2$ $(1\le j\le b)$, $q^{-4s}\xi_j^2$ $(b<j\le N)$, q^{-4s}/ξ_j^2 $(1\le j\le N)$, $q^{1-2s}w_a$, q^{-3-2s}/w_a $(a\in A)$. Here the integration contour $C^{(1)}$ is a closed curve such that w_a $(a\in A)$ encircles q^2/r in addition the same points $C^{(0)}$ does.

3.3 Identities between multiple integrals

From the spin-reversal properties in (2.37), (2.39), and (2.40), the following relation between form factors of local operators holds:

$$
F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r) = F^{(1-i,M,N)}_{-\epsilon_1,\cdots,-\epsilon_M,-\mu_1,\cdots,-\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;1/r). \tag{3.17}
$$

We should understand this spin-reversal property (3.17) as an analytic continuation of the parameter r. From the above (3.17), we obtain infinitely many relations between n-fold integrals which cannot be reduced to the relations between n-fold integrals of elliptic gamma functions summarized in [41]. Here we focus on two simple examples.

• First, from

$$F^{(0,M,N)}_{+,\cdots,+,+,\cdots,+}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r)=F^{(1,M,N)}_{-,\cdots,-,-,\cdots,-}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;1/r), \tag{3.18}$$

we obtain an identity for multiple integrals of infinite products. Here we consider the case $M=N$:

$$\begin{aligned}
&\prod_{j=1}^{M}\frac{(q^4r\zeta_j^2;q^4)_\infty(r/\zeta_j^2;q^4)_\infty(qr\xi_j^2;q^4)_\infty(qr/\xi_j^2;q^4)_\infty}{(q^2r\zeta_j^2;q^4)_\infty(q^2r/\zeta_j^2;q^4)_\infty(q^3r\xi_j^2;q^4)_\infty(r/q\xi_j^2;q^4)_\infty}\\
=\;& q^M(1-q^2)^Mq^{-3M(M+1)}(q^2;q^2)_\infty^{2M}\oint\cdots\oint_{C^{(0)}}\prod_{a=1}^{M}\frac{dw_a}{2\pi i}w_a\prod_{b=1}^{M}\frac{du_b}{2\pi i}\prod_{a=1}^{M}\frac{(1-ru_a/q^3)}{(1-rw_a/q^2)}\\
\times\;&\frac{\prod_{a=1}^{M}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty(u_a^2/q^2;q^4)_\infty(q^6/u_a^2;q^4)_\infty}{\prod_{a=1}^{M}\prod_{b=1}^{M}w_a^2(w_au_b/q^3;q^2)_\infty(q^3w_a/u_b;q^2)_\infty(u_b/qw_a;q^2)_\infty(q^5/w_au_b;q^2)_\infty}\\
\times\;&\frac{\prod_{1\le a<b\le M}w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a=1}^{M}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}}\\
\times\;&\frac{\prod_{1\le a<b\le M}u_a^2(u_au_b/q^4;q^2)_\infty(q^2u_a/u_b;q^2)_\infty(u_b/q^2u_a;q^2)_\infty(q^4/u_au_b;q^2)_\infty}{\prod_{a=1}^{M}\left\{\prod_{1\le j\le a}(\xi_j^2-q^{-4}u_a)\prod_{a\le j\le M}(u_a-q^2\xi_j^2)\right\}}\\
\times\;&\frac{\prod_{j=1}^{M}\prod_{b=1}^{M}u_b(q\zeta_j^2u_b;q^4)_\infty(q^3\zeta_j^2/u_b;q^4)_\infty(qu_b/\zeta_j^2;q^4)_\infty(q^3/\zeta_j^2u_b;q^4)_\infty}{\prod_{j=1}^{M}\prod_{a=1}^{M}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta_j^2;q^4)\infty(q^4/\zeta_j^2w_a;q^4)_\infty}\\
\times\;&\frac{\prod_{j=1}^{M}\prod_{a=1}^{M}w_a(q\xi_j^2w_a;q^4)_\infty(q^3\xi_j^2/w_q;q^4)_\infty(qw_a/\xi_j^2;q^4)_\infty(q^3/\xi_j^2w_a;q^4)_\infty}{\prod_{j=1}^{M}\prod_{b=1}^{M}(\xi_j^2u_b;q^4)_\infty(q^6\xi_j^2/u_b;q^4)_\infty(u_b/\xi_j^2;q^4)\infty(q^2/\xi_j^2u_b;q^4)_\infty}.
\end{aligned}\tag{3.19}$$

The number of integrals in the RHS is $(M+N)$ where the LHS is an infinite product without integral. Here the integration contour $C^{(0)}$ is given in the previous Section.

• Secondly, from

$$F^{(1,2m,0)}_{\underbrace{-,\cdots,-}_{m},\underbrace{+,\cdots,+}_{m}}(\zeta_1,\cdots,\zeta_{2m};r)=F^{(0,2m,0)}_{\underbrace{+,\cdots,+}_{m},\underbrace{-,\cdots,-}_{m}}(\zeta_1,\cdots,\zeta_{2m};1/r), \tag{3.20}$$

we obtain the following identity between multiple integrals of infinite products. The numbers of integrals on both sides are m.

$$\begin{aligned}
&\oint\cdots\oint_{C^{(0)}}\prod_{a=1}^{m}\frac{dw_a}{2\pi\sqrt{-1}}w_a\prod_{j=1}^{2m}\frac{(q^2\zeta_j^2/r;q^4)_\infty}{(q^4\zeta_j^2/r;q^4)_\infty}\prod_{a=1}^{m}\frac{1}{(1-w_a/q^2r)}\\
&\times\frac{\prod_{a=1}^{m}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty}{\prod_{j=1}^{2m}\prod_{a=1}^{m}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta\mathrm{J}^2;q^4)_\infty(q^4/\zeta_jw_a;q^4)_\infty}\\
&\times\frac{\prod_{1\le a<b\le m}w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a=1}^{m}\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le 2m}(w_a-q^4\zeta_j^2)}\\
=\;&(-q^3)^{-m^2+m}\prod_{j=m+1}^{2m}\zeta_j^2\oint\cdots\oint_{C^{(1)}}\prod_{a=m+1}^{2m}\frac{dw_a}{2\pi\sqrt{-1}}\prod_{j=1}^{2m}\frac{(1/r\zeta_j^2;q^4)_\infty}{(q^2/r\zeta_j^2;q^4)_\infty}\prod_{a=m+1}^{2m}\frac{1}{(1-q^2/rw_a)}
\end{aligned}$$

$$\times\frac{\prod_{a=m+1}^{2m}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty}{\prod_{j=1}^{2m}\prod_{a=m+1}^{2m}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta_j^2;q^4)_\infty(q^4/\zeta_jw_a;q^4)_\infty}$$
$$\times\frac{\prod_{m+1\le a<b\le 2m}w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a=m+1}^{2m}\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le 2m}(w_a-q^4\zeta_j^2)}. \tag{3.21}$$

Here the integration contour $C^{(0)}$ is given in the previous Section. In Appendix C, we give a direct proof for the simplest case where $\zeta_1=-q^{-1}\zeta$, $\zeta_2=\zeta$, and $m=1$ as a supporting argument :

$$F^{(1,2,0)}_{-,+}(-q^{-1}\zeta,\zeta;r)=F^{(0,2,0)}_{+,-}(-q^{-1}\zeta,\zeta;1/r). \tag{3.22}$$

3.4 Comparison with QISM

In this Section, upon specializations the previous expressions are compared with the known results obtained within the QISM [18]. In particular, let us consider the following integral representations of correlation functions :

$$\begin{aligned}
& g^mF^{(i,2m,0)}_{-\epsilon_1,\cdots,-\epsilon_m,\epsilon'_m,\cdots,\epsilon'_1}(\overbrace{-q^{-1}\zeta,\cdots,-q^{-1}\zeta}^{m},\overbrace{\zeta,\cdots,\zeta}^{m};r)\\
= \;& (-1)^{-\frac{1}{2}m^2-\frac{1}{2}m-\sum_{a\in A}a+(m+1)|B|}q^{\frac{1}{2}m^2+\frac{1}{2}m-2m|B|-\sum_{a\in A}a+\sum_{b\in B}b}\zeta^{2(m^2-m|B|-\sum_{a\in A}a)}\\
\times\;& \frac{(q^2;q^2)_\infty^{2m^2-m}}{(-q^2;q^2)_\infty^m}\left\{\frac{\Theta_{q^4}(\zeta^4)}{1-\zeta^4}\right\}^m\{(q^2\zeta^4;q^2)_\infty(q^2/\zeta^4;q^2)_\infty\}^{\frac{m(m-1)}{2}}(1-r\zeta^2)^m\oint\cdots\oint_{\widetilde{C}^{(i)}}\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}\prod_{b\in B}\frac{du_b}{2\pi\sqrt{-1}}\\
\times\;& \prod_{a\in A}\frac{w_a^{-2m+a}\Theta_{q^2}(w_a/q)\Theta_{q^2}(-w_a/q)(1-q^2/\zeta^2w_a)^m(1-w_a/\zeta^2)^{m-a}(1-q^2\zeta^2/w_a)^{a-1}}{\{\Theta_{q^2}(\zeta^2w_a)\Theta_{q^2}(w_a/\zeta^2)\}^m(1-rw_a/q^2)}\\
\times\;& \prod_{b\in B}\frac{u_b^{-m}\Theta_{q^2}(u_b/q)\Theta_{q^2}(-u_b/q)(1-q^2/\zeta^2u_b)^m(1-q^2\zeta^2/u_b)^{b-1}(1-q^4\zeta^2/u_b)^{m-b}}{\{\Theta_{q^2}(\zeta^2u_b)\Theta_{q^2}(u_b/\zeta^2)\}^m(1-ru_b/q^2)}\\
\times\;& \prod_{\substack{a<b\\a,b\in A}}\frac{w_a^2w_b^2\Theta_{q^2}(w_aw_b)\Theta_{q^2}(w_a/w_b)}{(1-q^2w_a/w_b)(1-q^4/w_aw_b)}\prod_{\substack{a>b\\a,b\in B}}\frac{u_a^2u_b^2\Theta_{q^2}(u_au_b)\Theta_{q^2}(u_a/u_b)}{(1-q^2u_a/u_b)(1-q^4/u_au_b)}\\
\times\;& \prod_{a\in A}\prod_{b\in B}\frac{w_a^2u_b^2\Theta_{q^2}(w_au_b)\Theta_{q^2}(w_a/u_b)}{(1-q^2w_a/u_b)(1-q^4/w_au_b)}.
\end{aligned} \tag{3.23}$$

In this Section we set $A=\{1\le j\le n|\epsilon_j=-\}$, $B=\{1\le j\le|\epsilon'_j=+\}$. Here the integration contour $\widetilde{C}^{(0)}$ is a closed curve that satisfies the following conditions for $s=0,1,2,\cdots$. The $w_a(a\in A)$ encircles $q^{2+2s}\zeta^2$, q^{4+2s}/ζ^2, but not $q^{-2-2s}\zeta^2$, q^{-2s}/ζ^2, and q^2/r. The $u_b(b\in B)$ encircles $q^{4+2s}\zeta^2$, q^{4+2s}/ζ^2, but not $q^{-2s}\zeta^2$, q^{-2s}/ζ^2, and q^2/r. The integral contour $\widetilde{C}^{(1)}$ is a closed curve such that $w_a(a\in A)$ and $u_b(b\in B)$ encircle q^2/r in addition the same points $\widetilde{C}^{(0)}$ does.

In what follows we study integral representations upon the specialization of the spectral parameter $\zeta=1$. We change the variables as follows.

$$q=e^{-\eta}\ (\eta>0),\quad w_a/q=e^{2\sqrt{-1}\alpha_a},\quad u_b/q=e^{2\sqrt{-1}\delta_b},\quad r=e^{2\xi}\ (\xi\in\mathbf{R}). \tag{3.24}$$

We use the following notation of the elliptic theta functions in [42].

$$\vartheta_1(u,q)\;=\;2q^{\frac{1}{4}}(q^2;q^2)_\infty\sin u\prod_{n=1}^\infty(1-2q^{2n}\cos 2u+q^{4n}), \tag{3.25}$$

$$\vartheta_2(u,q) = 2q^{\frac{1}{4}}(q^2;q^2)_\infty \cos u \prod_{n=1}^{\infty}(1+2q^{2n}\cos 2u + q^{4n}), \tag{3.26}$$

$$\vartheta_3(u,q) = (q^2;q^2)_\infty \prod_{n=1}^{\infty}(1+2q^{2n-1}\cos 2u + q^{4n-2}), \tag{3.27}$$

$$\vartheta_4(u,q) = (q^2;q^2)_\infty \prod_{n=1}^{\infty}(1-2q^{2n-1}\cos 2u + q^{4n-2}). \tag{3.28}$$

The theta function $\Theta_p(w) = (p;p)_\infty (w;p)_\infty (p/w;p)_\infty$ is related to $\vartheta_j(\alpha,q)$ as follows.

$$\Theta_{q^2}(w) = \vartheta_4(\alpha,q), \quad \Theta_{q^2}(w/q) = \frac{\sqrt{-1}q^{\frac{1}{4}}}{\sqrt{w}}\vartheta_1(\alpha,q), \quad \Theta_{q^2}(-w/q) = \frac{q^{\frac{1}{4}}}{\sqrt{w}}\vartheta_2(\alpha,q). \tag{3.29}$$

Here we have set $w/q = e^{2\sqrt{-1}\alpha}$.

• The following is the main result of this Section. For $|A|+|B| = m$, we have the following integral representation of correlation functions :

$$\begin{aligned}
&\frac{{}_B\langle i|(E_{\epsilon_1,\epsilon_1'})_1(E_{\epsilon_2,\epsilon_2'})_2\cdots(E_{\epsilon_m,\epsilon_m'})_m|i\rangle_B}{{}_B\langle i|i\rangle_B} \\
= \;& g^m F^{(i,2m,0)}_{-\epsilon_1,\cdots,-\epsilon_m,\epsilon_m',\cdots,\epsilon_1'}(\underbrace{-q^{-1},\cdots,-q^{-1}}_{m},\underbrace{1,\cdots,1}_{m};r) \\
= \;& 2^{m(m+1)}\sinh^m(\xi)\times(q^2;q^2)_\infty^{m(3m+1)}(-q^2;q^2)_\infty^{2m} \\
\times \;& (-1)^{\frac{m(m-1)}{2}+m(|B|-|A|)}q^{-\frac{m(m-1)}{4}+m(m-\frac{1}{2})}\int\cdots\int_{\widetilde{C}^{(i)}}\prod_{a\in A}\frac{d\alpha_a}{2\pi}\sqrt{-1}\prod_{b\in B}\frac{d\delta_b}{2\pi\sqrt{-1}} \\
\times \;& \prod_{a\in A}\frac{\vartheta_1(\alpha_a,q)\vartheta_2(\alpha_a,q)\;\sin^{m+a-1}\left(\alpha_a+\frac{\sqrt{-1}}{2}\eta\right)\sin^{m-a}(\alpha_a)}{\sin\left(\alpha_a+\frac{\sqrt{-1}}{2}\eta+\sqrt{-1}\xi\right)\;\vartheta_4(\alpha_a,q)^{2n}} \\
\times \;& \prod_{b\in B}\frac{\vartheta_1(\delta_b,q)\vartheta_2(\delta_b,q)\;\sin^{m+b-1}\left(\delta_b+\frac{\sqrt{-1}}{2}\eta\right)\sin^{m-b}\left(\delta_b+\frac{3\sqrt{-1}}{2}\eta\right)}{\sin\left(\delta_b+\frac{\sqrt{-1}}{2}\eta+\sqrt{-1}\xi\right)\;\vartheta_4(\delta_b,q)^{2m}} \\
\times \;& \prod_{\substack{a<b\\a,b\in A}}\frac{\vartheta_1(\alpha_a+\alpha_b,q)\vartheta_1(\alpha_a-\alpha_b,q)}{\sin(\alpha_a+\alpha_b+\sqrt{-1}\eta)\sin(\alpha_a-\alpha_b-\sqrt{-1}\eta)}\prod_{\substack{a>b\\a,b\in B}}\frac{\vartheta_1(\delta_a+\delta_b,q)\vartheta_1(\delta_a-\delta_b,q)}{\sin(\delta_a+\delta_b+\sqrt{-1}\eta)\sin(\delta_a-\delta_b-\sqrt{-1}\eta)} \\
\times \;& \prod_{a\in A}\prod_{b\in B}\frac{\vartheta_1(\alpha_a+\delta_b,q)\vartheta_1(\alpha_a-\delta_b,q)}{\sin(\alpha_a+\delta_b+\sqrt{-1}\eta)\sin(\alpha_a-\delta_b-\sqrt{-1}\eta)}.
\end{aligned} \tag{3.30}$$

Note that we obtain the same formula as the one of Lyon group [18] up to a constant $(-1)^{m(|B|-|A|)}q^{m(m-\frac{1}{2})}$.

$$\text{“(3.30) of this paper”} = (-1)^{m(|B|-|A|)}q^{m(m-\frac{1}{2})}\times\text{“(6.18) of [18]”}. \tag{3.31}$$

4 Form factors for a triangular boundary

In this Section, we study the form factors of local operators for a *triangular* boundary. Two integral representations of form factors of local operators are proposed using bosonization.

4.1 Definition

In this Section we study the vacuum expectation values of the q-vertex operators with natural numbers M, N with $M = N \ (mod.2)$:

$$\begin{aligned} & F^{(i,\pm,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s) \\ = \ & \frac{{}_B\langle i;\pm|\Phi^{(i,i+1)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(i+M-1,i+M)}_{\epsilon_M}(\zeta_M)\Psi^{*(i+N,i+N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(i+1,i)}_{\mu_N}(\xi_N)|\pm;i\rangle_B}{{}_B\langle i;\pm|\pm;i\rangle_B}. \end{aligned} \tag{4.1}$$

Here the number i in the suffix $(i, i+1)$ of q-vertex operators $\Phi^{(i,i+1)}_{\epsilon}(\zeta)$ should be understood modulo 2. Let us consider the form factors of local operators with $M = 2m$ and N : even given by

$$\begin{aligned} & {}_B\langle i;\pm|(\mathcal{E}_{\epsilon_m\epsilon'_m})_m(\mathcal{E}_{\epsilon_{m-1}\epsilon'_{m-1}})_{m-1}\cdots(\mathcal{E}_{\epsilon_1\epsilon'_1})_1\Psi^{*(i+N,i+N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(i+1,i)}_{\mu_N}(\xi_N)|\pm;i\rangle_B \\ & = g^m \times \ {}_B\langle i;\pm|\pm;i\rangle_B \times F^{(\pm,i,M,N)}_{-\epsilon_1,\cdots,-\epsilon_m,\epsilon'_m,\cdots,\epsilon'_1,\mu_1,\cdots,\mu_N}(\underbrace{-q^{-1},\cdots,-q^{-1}}_{m},\underbrace{1,\cdots,1}_{m},\xi_1,\cdots,\xi_N;r,s). \end{aligned} \tag{4.2}$$

In what follows we call both (4.1) and (4.2) the form factors of local operators. The form factors are calculated using bosonizations.

4.2 First integral representation

In this Section we focus our attention on the half-infinite XXZ spin chain with a upper-triangular boundary. We calculate the form factors with $M = N \ (mod.2)$:

$$F^{(+,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s). \tag{4.3}$$

Calculations in this Section are similar to those for diagonal boundary conditions. In what follows we set $A = \{1 \le a \le M | \epsilon_a = +\}$, $B = \{1 \le b \le N | \mu_b = +\}$, and

$$L = |A| - |B| + \frac{1}{2}(N - M). \tag{4.4}$$

In the following, we understand $L = 0$ as a sufficient condition such that the expressions for triangular boundary conditions coincide with those for diagonal boundary conditions.

The following bosonizations of the Chevalley generators are used:

$$f_0 = q^{\partial}\oint\frac{dv}{2\pi\sqrt{-1}}X^+(v), \quad e_1q^{-h_1} = \oint\frac{dv}{2\pi\sqrt{-1}}X^+(v)q^{-\partial}. \tag{4.5}$$

The bosonic parts of the following operators :

$$\begin{aligned} & \exp_q(-sf_0)\cdot\Phi^{(0,1)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(M-1,M)}_{\epsilon_M}(\zeta_M)\cdot\Psi^{*(N,N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(1,0)}_{\mu_N}(\xi_N)\cdot\exp_{q^{-1}}(sf_0), \\ & \exp_{q^{-1}}\left(-\frac{s}{rq}e_1q^{-h_1}\right)\cdot\Phi^{(1,0)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(M,M+1)}_{\epsilon_M}(\zeta_M)\cdot\Psi^{*(N+1,N)}_{\mu_1}(\xi_1)\cdots\Psi^{*(0,1)}_{\mu_N}(\xi_N)\cdot\exp_q\left(\frac{s}{rq}e_1q^{-h_1}\right), \end{aligned} \tag{4.6}$$

are given by

$$:\prod_{j=1}^{M}\Phi^{(i+j-1,i+j)}_{-}(\zeta_j)\prod_{j=1}^{N}\Psi^{*(i+N-j+1,i+N-j)}_{-}(\xi_j)\prod_{a\in A}X^-(w_a)\prod_{a=1}^{n}X^+(v_a)\prod_{a\in B}X^+(u_a):q^{(1-2i)n\partial}, \tag{4.7}$$

for $n=0,1,2,\cdots$. Here the parameter n comes from the expansion of the q-exponential. The zero-mode of the above operator is $e^{\alpha(n-|A|+|B|+\frac{1}{2}(M-N))}$, hence the vacuum expectation value becomes non-zero i.e.:

$$ {}_B\langle +;i|:\prod_{j=1}^{M}\Phi_{-}^{(i+j-1,i+j)}(\zeta_j)\prod_{j=1}^{N}\Psi_{-}^{*(i+N-j+1,i+N-j)}(\xi_j)\prod_{a\in A}X^{-}(w_a)\prod_{a=1}^{n}X^{+}(v_a)\prod_{a\in B}X^{+}(u_a):|i;+\rangle_B\neq 0, \tag{4.8}$$

if and only if $L=n\geq 0$.

For $L=|A|-|B|+\frac{1}{2}(N-M)\geq 0$, we have the vacuum expectation values of the q-vertex operators takes the form:

$$\begin{aligned}
& {}_B\langle i;+|\Phi_{\epsilon_1}^{(i,i+1)}(\zeta_1)\cdots\Phi_{\epsilon_M}^{(i+M-1,i+M)}(\zeta_M)\Psi_{\mu_1}^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_{\mu_N}^{*(i+1,i)}(\xi_N)|+;i\rangle_B \\
= \;& (qs)^{|A|-|B|+\frac{1}{2}(N-M)}(rq^2)^{(|B|-|A|+\frac{1}{2}(M-N))i}q^{|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}(1-q^2)^{|A|+|B|} \\
\times \;& \sum_{\substack{l,m\geq 0\\ l+m=|A|-|B|+\frac{1}{2}(-M+N)}}\frac{(-1)^{l+mN}q^{\frac{-l(l+1)+m(m+1)}{2}(1-2i)-3Nm}}{[l]_q![m]_q!}\prod_{j=1}^{M}\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i}\prod_{j=1}^{N}\xi_j^{\frac{1+\mu_j}{2}+N-j+1-i} \\
\times \;& \prod_{1\leq j<k\leq M}\frac{(q^2\zeta_k^2/\zeta_j^2;q^4)_\infty}{(q^4\zeta_k^2/\zeta_j^2;q^4)_\infty}\prod_{1\leq j<k\leq N}\frac{(\xi_k^2/\xi_j^2;q^4)_\infty}{(q^2\xi_k^2/\xi_j^2;q^4)_\infty}\prod_{j=1}^{M}\prod_{k=1}^{N}\frac{(q^3\xi_k^2/\zeta_j^2;q^4)_\infty}{(q\xi_k^2/\zeta_j^2;q^4)_\infty} \\
\times \;& \oint\cdots\oint\prod_{a=1}^{l+m}\frac{dv_a}{2\pi\sqrt{-1}}v_a^{2i-1}\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a^{1-i}\prod_{a\in B}\frac{du_a}{2\pi\sqrt{-1}}u_a^{i} \\
\times \;& \frac{\prod_{a\in A}\prod_{j=1}^{N}(w_a-q^3\xi_j^2)\prod_{a\in B}\prod_{j=1}^{M}(u_a-q^3\zeta_j^2)\prod_{a=1}^{l+m}\prod_{j=1}^{M}(v_a-q^3\zeta_j^2)}{\prod_{a\in A}\left\{\prod_{1\leq j\leq a}(\zeta_j^2-q^{-2}w_a)\prod_{a\leq j\leq M}(w_a-q^4\zeta_j^2)\right\}\prod_{a\in B}\left\{\prod_{1\leq j\leq a}(\xi_j^2-q^{-4}u_a)\prod_{a\leq j\leq N}(u_a-q^2\xi_j^2)\right\}} \\
\times \;& \frac{\prod_{\substack{a,b\in A\\ a<b}}(w_a-w_b)(w_a-q^2w_b)\prod_{\substack{a,b\in B\\ a<b}}(u_a-u_b)(u_a-q^{-2}u_b)\prod_{1\leq a<b\leq l+m}(v_a-v_b)(v_a-q^{-2}v_b)}{\prod_{j=1}^{N}\left\{\prod_{a=1}^{l}(v_a-q^2\xi_j^2)\prod_{a=l+1}^{l+m}(\xi_j^2-q^{-4}v_a)\right\}} \\
\times \;& \frac{\prod_{b\in B}\left\{\prod_{a=1}^{l}(v_a-u_b)(v_a-q^{-2}u_b)\prod_{a=l+1}^{l+m}(u_b-v_a)(u_b-q^{-2}v_a)\right\}}{\prod_{a\in A}\left\{\prod_{b\in B}(w_a-qu_b)(w_a-q^{-1}u_b)\prod_{b=1}^{l+m}(w_a-qv_b)(w_a-q^{-1}v_b)\right\}} \\
\times \;& {}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^{-}(w_a)+\sum_{a\in B}R^{+}(u_a)+\sum_{a=1}^{l+m}R^{+}(v_a)} \\
\times \;& e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^{-}(w_a)+\sum_{a\in B}S^{+}(u_a)+\sum_{a=1}^{l+m}S^{+}(v_a)}|i\rangle_B.
\end{aligned}\tag{4.9}$$

Using (3.15), we calculate the following vacuum expectation value :

$$\begin{aligned}
& {}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^{-}(w_a)+\sum_{a\in B}R^{+}(u_a)+\sum_{a=1}^{L}R^{+}(v_a)} \\
\times \;& e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^{-}(w_a)+\sum_{a\in B}S^{+}(u_a)+\sum_{a=1}^{L}S^{+}(v_a)}|i\rangle_B \\
= \;& \langle i|e^{G^{(i)}}e^{\sum_{n=1}^{\infty}a_{-n}X_n}e^{-\sum_{n=1}^{\infty}a_nY_n}e^{F^{(i)}}|i\rangle.
\end{aligned}\tag{4.10}$$

Here we have defined

$$X_n = \frac{q^{7n/2}}{[2n]_q}\sum_{j=1}^{M}\zeta_j^{2n}-\frac{q^{5n/2}}{[2n]_q}\sum_{j=1}^{N}\xi_j^{2n}-\frac{q^{n/2}}{[n]_q}\sum_{a\in A}w_a^n+\frac{q^{-n/2}}{[n]_q}\sum_{a\in B}u_a^n+\frac{q^{-n/2}}{[n]_q}\sum_{a=1}^{L}v_a^n, \tag{4.11}$$

$$Y_n = \frac{q^{-5n/2}}{[2n]_q}\sum_{j=1}^{M}\zeta_j^{-2n}-\frac{q^{-7n/2}}{[2n]_q}\sum_{j=1}^{N}\xi_j^{-2n}-\frac{q^{n/2}}{[n]_q}\sum_{a\in A}w_a^{-n}+\frac{q^{-n/2}}{[n]_q}\sum_{a\in B}u_a^{-n}+\frac{q^{-n/2}}{[n]_q}\sum_{a=1}^{L}v_a^{-n}. \tag{4.12}$$

- The followings is the main results of this Section. For $L=|A|-|B|+\frac{1}{2}(N-M)<0$, we have

$$F^{(+,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)=0. \tag{4.13}$$

For $L=|A|-|B|+\frac{1}{2}(N-M)=0$, the form factors for triangular boundary conditions coincide with those for diagonal boundary conditions :

$$F^{(+,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)=F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r). \tag{4.14}$$

For $L=|A|-|B|+\frac{1}{2}(N-M)>0$, the following integral representations of the form factors of local operators are obtained:

$$\begin{aligned}
&F^{(+,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)\\
=\;&(qs)^L(rq^2)^{-Li}q^{|B|}(1-q^2)^{|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}\\
\times\;&\left(\frac{\{q^6\}_\infty}{\{q^8\}_\infty}\right)^M\left(\frac{\{q^4\}_\infty}{\{q^6\}_\infty}\right)^N\{(q^2;q^2)_\infty\}^{2|A|+\frac{1}{2}(-M+N)}\prod_{j=1}^M\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i}\prod_{j=1}^N\xi_j^{\frac{\mu_j+1}{2}+N-j+1-i}\\
\times\;&\prod_{1\le j<k\le M}\frac{\{q^6\zeta_j^2\zeta_k^2\}_\infty\{q^2/\zeta_j^2\zeta_k^2\}_\infty\{q^6\zeta_j^2/\zeta_k^2\}_\infty\{q^2\zeta_k^2/\zeta_j^2\}_\infty}{\{q^8\zeta_j^2\zeta_k^2\}_\infty\{q^4/\zeta_j^2\zeta_k^2\}_\infty\{q^8\zeta_j^2/\zeta_k^2\}_\infty\{q^4\zeta_k^2/\zeta_j^2\}_\infty}\\
\times\;&\prod_{1\le j<k\le N}\frac{\{q^4\xi_j^2\xi_k^2\}_\infty\{1/\xi_j^2\xi_k^2\}_\infty\{q^4\xi_j^2/\xi_k^2\}_\infty\{\xi_k^2/\xi_j^2\}_\infty}{\{q^6\xi_j^2\xi_k^2\}_\infty\{q^2/\xi_j^2\xi_k^2\}_\infty\{q^6\xi_j^2/\xi_k^2\}_\infty\{q^2\xi_k^2/\xi_j^2\}_\infty}\\
\times\;&\prod_{j=1}^M\prod_{k=1}^N\frac{\{q^7\zeta_j^2\xi_k^2\}_\infty\{q^3/\zeta_j^2\xi_k^2\}_\infty\{q^7\zeta_j^2/\xi_k^2\}_\infty\{q^3\xi_k^2/\zeta_j^2\}_\infty}{\{q^5\zeta_j^2\xi_k^2\}_\infty\{q/\zeta_j^2\xi_k^2\}_\infty\{q^5\zeta_j^2/\xi_k^2\}_\infty\{q\xi_k^2/\zeta_j^2\}_\infty}\\
\times\;&\prod_{j=1}^M\frac{[q^{10}\zeta_j^4]_\infty[q^{14}\zeta_j^4]_\infty[q^{10}/\zeta_j^4]_\infty[q^6/\zeta_j^4]_\infty}{[q^{12}\zeta_j^4]_\infty[q^{16}\zeta_j^4]_\infty[q^{12}/\zeta_j^4]_\infty[q^8/\zeta_j^4]_\infty}\prod_{j=1}^N\frac{[q^4\xi_j^4]_\infty[q^8\xi_j^4]_\infty[1/\xi_j^4]_\infty[q^4/\xi_j^4]_\infty}{[q^6\xi_j^4]_\infty[q^{10}\xi_j^4]_\infty[q^2/\xi_j^4]_\infty[q^6/\xi_j^4]_\infty}\\
\times\;&\sum_{\substack{l,m\ge0\\ l+m=L}}\frac{(-1)^{l+mN}q^{\frac{l(l+1)-m(m+1)}{2}(2i-1)-3Nm}}{[l]_q![m]_q!}\oint\cdots\oint_{C_l^{(+,i)}}\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a^{1-i}\prod_{b\in B}\frac{du_b}{2\pi\sqrt{-1}}u_b^i\prod_{c=1}^L\frac{dv_c}{2\pi\sqrt{-1}}v_c^{2i-1}\\
&\prod_{a\in A}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty\prod_{a\in B}(u_a^2/q^2;q^4)_\infty(q^6/u_a^2;q^4)_\infty\prod_{a=1}^L(v_a^2/q^2;q^4)_\infty(q^6/v_a^2;q^4)_\infty\\
\times\;&\frac{\prod_{1\le a<b\le L}v_a^2(v_av_b/q^4;q^2)_\infty(q^2v_a/v_b;q^2)_\infty(v_b/q^2v_a;q^2)_\infty(q^4/v_av_b;q^2)_\infty}{\prod_{a\in A}\prod_{b\in B}w_a^2(w_au_b/q^3;q^2)_\infty(q^3w_a/u_b;q^2)_\infty(u_b/qw_a;q^2)_\infty(q^5/w_au_b;q^2)_\infty}\\
\times\;&\frac{\prod_{a=1}^L\prod_{b\in B}(v_au_b/q^4;q^2)_\infty(q^2v_a/u_b;q^2)_\infty(q^2u_b/v_a;q^2)_\infty(q^4/v_au_b;q^2)_\infty}{\prod_{a=1}^L\prod_{b\in A}v_a^2(v_aw_b/q^3;q^2)_\infty(q^3v_a/w_b;q^2)_\infty(w_b/qv_a;q^2)_\infty(q^5/v_aw_b;q^2)_\infty}\\
\times\;&\frac{\prod_{\substack{a,b\in A\\ a<b}}w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a\in A}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}}\\
\times\;&\frac{\prod_{\substack{a,b\in B\\ a<b}}u_a^2(u_au_b/q^4;q^2)_\infty(q^2u_a/u_b;q^2)_\infty(u_b/q^2u_a;q^2)_\infty(q^4/u_au_b;q^2)_\infty}{\prod_{a\in B}\left\{\prod_{1\le j\le a}(\xi_j^2-q^{-4}u_a)\prod_{a\le j\le M}(u_a-q^2\xi_j^2)\right\}}\\
\times\;&\frac{\prod_{j=1}^M\prod_{a\in B}u_a(q\zeta_j^2u_a;q^4)_\infty(q^3\zeta_j^2/u_a;q^4)_\infty(qu_a/\zeta_j^2;q^4)_\infty(q^3/\zeta_j^2u_a;q^4)_\infty}{\prod_{j=1}^M\prod_{a\in A}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta_j^2;q^4)_\infty(q^4/\zeta_j^2w_a;q^4)_\infty}
\end{aligned}$$

$$
\begin{aligned}
&\times \frac{\prod_{j=1}^{N}\prod_{a\in A} w_a(q\xi_j^2 w_a;q^4)_\infty (q^3\xi_j^2/w_a;q^4)_\infty (qw_a/\xi_j^2;q^4)_\infty (q^3/\xi_j^2 w_a;q^4)_\infty}{\prod_{j=1}^{N}\prod_{a\in B}(\xi_j^2 u_a;q^4)_\infty (q^6\xi_j^2/u_a;q^4)_\infty (u_a/\xi_j^2;q^4)_\infty (q^2/\xi_j^2 u_a;q^4)_\infty} \\
&\times \frac{\prod_{j=1}^{M}\prod_{a=1}^{L} v_a(q\zeta_j^2 v_a;q^4)_\infty (q^3\zeta_j^2/v_a;q^4)_\infty (qv_a/\zeta_j^2;q^4)_\infty (q^3/\zeta_j^2 v_a;q^4)_\infty}{\prod_{j=1}^{N}\prod_{a=1}^{L}(\xi_j^2 v_a;q^4)_\infty (q^6\xi_j^2/v_a;q^4)_\infty (v_a/\xi_j^2;q^4)_\infty (q^2/\xi_j^2 v_a;q^4)_\infty} \\
&\times \frac{\prod_{b\in B}\left\{\prod_{a=1}^{l}(v_a-u_b)(v_a-q^{-2}u_b)\prod_{a=l+1}^{L}(u_b-v_a)(u_b-q^{-2}v_a)\right\}}{\prod_{j=1}^{N}\left\{\prod_{a=1}^{l}(v_a-q^2\xi_j^2)\prod_{a=l+1}^{L}(\xi_j^2-q^{-4}v_a)\right\}} \\
&\times \begin{cases} \prod_{j=1}^{M}\frac{(q^2 r\zeta_j^2;q^4)_\infty}{(q^4 r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q^3 r\xi_j^2;q^4)_\infty}{(qr\xi_j^2;q^4)_\infty}\frac{\prod_{a\in B}(1-ru_a/q^3)\prod_{a=1}^{L}(1-rv_a/q^3)}{\prod_{a\in A}(1-rw_a/q^2)} & (i=0), \\ \prod_{j=1}^{M}\frac{(1/r\zeta_j^2;q^4)_\infty}{(q^2/r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q/r\xi_j^2;q^4)_\infty}{(1/qr\xi_j^2;q^4)_\infty}\frac{\prod_{a\in B}(1-q/ru_a)\prod_{a=1}^{L}(1-q/rv_a)}{\prod_{a\in A}(1-q^2/rw_a)} & (i=1). \end{cases}
\end{aligned}
\tag{4.15}
$$

Here the integration contour $C_l^{(+,0)}$ $(1\le l\le L)$ is a closed curve that satisfies the following conditions for $s=0,1,2,\cdots$. The w_a $(a\in A)$ encircles $q^{8+4s}\zeta_j^2$ $(1\le j<a)$, $q^{4+4s}\zeta_j^2$ $(a\le j\le M)$, q^{4+4s}/ζ_j^2 $(1\le j\le M)$, $q^{-1+2s}u_b$, q^{5+2s}/u_b $(b\in B)$, $q^{3+2s}v_c$ $(1\le c\le l)$, $q^{-1+2s}v_c$ $(l<c\le L)$, q^{5+2s}/v_c $(1\le c\le L)$, but not $q^{2-4s}\zeta_j^2$ $(1\le j\le a)$, $q^{-2-4s}\zeta_j^2$ $(a<j\le M)$, q^{-2-4s}/ζ_j^2 $(1\le j\le M)$, $q^{-3-2s}u_b$, q^{3-2s}/u_b $(b\in B)$, q^2/r, $q^{1-2s}v_c$ $(1\le c\le l)$, $q^{-3-2s}v_c$ $(l<c\le L)$, q^{3-2s}/v_c $(1\le c\le L)$. The u_b $(b\in B)$ encircles $q^{6+4s}\xi_j^2$ $(1\le j<b)$, $q^{2+4s}\xi_j^2$ $(b\le j\le N)$, q^{2+4s}/ξ_j^2 $(1\le j\le N)$, $q^{3+2s}w_a$, q^{5+2s}/w_a $(a\in B)$, but not $q^{4-4s}\xi_j^2$ $(1\le j\le b)$, $q^{-4s}\xi_j^2$ $(b<j\le N)$, q^{-4s}/ξ_j^2 $(1\le j\le N)$, $q^{1-2s}w_a$, q^{-3-2s}/w_a $(a\in A)$. The v_c $(1\le c\le l)$ encircles $q^{2+4s}\xi_j^2$, q^{2+4s}/ξ_j^2 $(1\le j\le N)$, $q^{-1+2s}w_a$, q^{5+2s}/w_a $(a\in A)$, but not $q^{-4s}\xi_j^2$, q^{-4s}/ξ_j^2 $(1\le j\le N)$, $q^{-3-2s}w_a$, q^{3-2s}/w_a $(a\in A)$. The v_c $(l<c\le L)$ encircles $q^{6+4s}\xi_j^2$, q^{2+4s}/ξ_j^2 $(1\le j\le N)$, $q^{3+2s}w_a$, q^{5+2s}/w_a $(a\in A)$, but not $q^{4-4s}\xi_j^2$, q^{-4s}/ξ_j^2 $(1\le j\le N)$, $q^{1-2s}w_a$, q^{3-2s}/w_a $(a\in A)$. Here the integration contour $C_l^{(+,1)}$ $(1\le l\le L)$ is a closed curve such that w_a $(a\in A)$ encircles q^2/r in addition the same points $C_l^{(+,0)}$ does.

4.3 Second integral representation

In this Section we focus our attention on the half-infinite XXZ spin chain with a lower-triangular boundary. Let us consider the form factors with $M=N$ $(mod.2)$:

$$
F^{(-,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s). \tag{4.16}
$$

In what follows we use $A=\{1\le a\le M|\epsilon_a=+\}$, $B=\{1\le b\le N|\mu_b=+\}$, and

$$
K=|B|-|A|+\frac{1}{2}(M-N). \tag{4.17}
$$

In the following, we understand $K=0$ as a sufficient condition such that the expressions for triangular boundary conditions coincide with those for diagonal boundary conditions. The following bosonizations of the Chevalley generators are used:

$$
f_1=\oint\frac{dv}{2\pi\sqrt{-1}}X^-(v),\quad e_0q^{-h_0}=q^{-1}\oint\frac{dv}{2\pi\sqrt{-1}}vX^+(v). \tag{4.18}
$$

The bosonic parts of the following operators :

$$
\begin{aligned}
&\exp_{q^{-1}}\left(\frac{s}{q}e_0q^{-h_0}\right)\Phi^{(0,1)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(M-1,M)}_{\epsilon_M}(\zeta_M)\Psi^{*(N,N-1)}_{\mu_1}(\xi_1)\cdots\Psi^{*(1,0)}_{\mu_N}(\xi_N)\exp_q\left(-\frac{s}{q}e_0q^{-h_0}\right), \\
&\exp_q\left(\frac{s}{r}f_1\right)\Phi^{(1,0)}_{\epsilon_1}(\zeta_1)\cdots\Phi^{(M,M+1)}_{\epsilon_M}(\zeta_M)\Psi^{*(N+1,N)}_{\mu_1}(\xi_1)\cdots\Psi^{*(0,1)}_{\mu_{2N}}(\xi_N)\exp_{q^{-1}}\left(-\frac{s}{r}f_1\right),
\end{aligned}
\tag{4.19}
$$

are given by

$$:\prod_{j=1}^{M}\Phi_{-}^{(i+j-1,i+j)}(\zeta_j)\prod_{j=1}^{N}\Psi_{-}^{*(i+N-j+1,i+N-j)}(\xi_j)\prod_{a\in A}X^{-}(w_a)\prod_{a\in B}X^{+}(u_a)\prod_{a=1}^{n}X^{-}(v_a):, \tag{4.20}$$

for $n=0,1,2,\cdots$. Here the parameter n comes from the q-exponential. The zero-mode of the above operator is $e^{\alpha(-n-|A|+|B|+\frac{1}{2}(M-N))}$, hence the vacuum expectation value becomes non-zero :

$$ {}_B\langle -;i|:\prod_{j=1}^{M}\Phi_{-}^{(i+j-1,i+j)}(\zeta_j)\prod_{j=1}^{N}\Psi_{-}^{*(i+N-j+1,i+N-j)}(\xi_j)\prod_{a\in A}X^{-}(w_a)\prod_{a\in B}X^{+}(u_a)\prod_{a=1}^{n}X^{-}(v_a):|-;i\rangle_B\neq 0, \tag{4.21}$$

if and only if $K=n\geq 0$.

For $K=|B|-|A|+\frac{1}{2}(M-N)\geq 0$, the vacuum expectation values of the q-vertex operators read:

$$\begin{aligned}
&{}_B\langle i;-|\Phi_{\epsilon_1}^{(i,i+1)}(\zeta_1)\cdots\Phi_{\epsilon_M}^{(i+M-1,i+M)}(\zeta_M)\Psi_{\mu_1}^{*(i+N,i+N-1)}(\xi_1)\cdots\Psi_{\mu_N}^{*(1-i,i)}(\xi_N)|-;i\rangle_B\\
=\;&(s/q^2)^{\frac{1}{2}(M-N)-|A|+|B|}(q^2/r)^{(\frac{1}{2}(M-N)-|A|+|B|)i}q^{|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}(1-q^2)^{|A|+|B|}\\
\times\;&\sum_{\substack{l,m\geq 0\\ l+m=\frac{1}{2}(M-N)-|A|+|B|}}\frac{(-1)^{m(M+1)}q^{\frac{l(l-1)-m(m-1)}{2}(2i-1)-3Mm}}{[l]_q![m]_q!}\prod_{j=1}^{M}\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i}\prod_{j=1}^{N}\xi_j^{\frac{1+\mu_j}{2}+N-j+1-i}\\
\times\;&\prod_{1\leq j<k\leq M}\frac{(q^2\zeta_k^2/\zeta_j^2;q^4)_\infty}{(q^4\zeta_k^2/\zeta_j^2;q^4)_\infty}\prod_{1\leq j<k\leq N}\frac{(\xi_k^2/\xi_j^2;q^4)_\infty}{(q^2\xi_k^2/\xi_j^2;q^4)_\infty}\prod_{j=1}^{M}\prod_{k=1}^{N}\frac{(q^3\xi_k^2/\zeta_j^2;q^4)_\infty}{(q\xi_k^2/\zeta_j^2;q^4)_\infty}\\
\times\;&\oint\cdots\oint\prod_{a=1}^{l+m}\frac{dv_a}{2\pi\sqrt{-1}}v_a^{1-2i}\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a^{1-i}\prod_{a\in B}\frac{du_a}{2\pi\sqrt{-1}}u_a^{i}\\
\times\;&\frac{\prod_{a\in A}\prod_{j=1}^{N}(w_a-q^3\xi_j^2)\prod_{a\in B}\prod_{j=1}^{M}(u_a-q^3\zeta_j^2)\prod_{a=1}^{l+m}\prod_{j=1}^{N}(v_a-q^3\xi_j^2)}{\prod_{a\in A}\left\{\prod_{1\leq j\leq a}(\zeta_j^2-q^{-2}w_a)\prod_{a\leq j\leq M}(w_a-q^4\zeta_j^2)\right\}\prod_{a\in B}\left\{\prod_{1\leq j\leq a}(\xi_j^2-q^{-4}u_a)\prod_{a\leq j\leq N}(u_a-q^2\xi_j^2)\right\}}\\
\times\;&\frac{\prod_{\substack{a,b\in A\\ a<b}}(w_a-w_b)(w_a-q^2w_b)\prod_{\substack{a,b\in B\\ a<b}}(u_a-u_b)(u_a-q^{-2}u_b)\prod_{1\leq a<b\leq l+m}(v_a-v_b)(v_a-q^2v_b)}{\prod_{j=1}^{M}\left\{\prod_{a=1}^{l}(v_a-q^4\zeta_j^2)\prod_{a=l+1}^{l+m}(\zeta_j^2-q^{-2}v_a)\right\}}\\
\times\;&\frac{\prod_{a\in A}\left\{\prod_{b=1}^{l}(v_b-w_a)(v_b-q^2w_a)\prod_{b=l+1}^{l+m}(w_a-v_b)(w_a-q^2v_b)\right\}}{\prod_{b\in B}\left\{\prod_{a\in A}(w_a-qu_b)(w_a-q^{-1}u_b)\prod_{a=1}^{l+m}(v_a-qu_b)(v_a-q^{-1}u_b)\right\}}\\
\times\;&{}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^{-}(w_a)+\sum_{a\in B}R^{+}(u_a)+\sum_{a=1}^{l+m}R^{-}(v_a)}\\
\times\;&e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^{-}(w_a)+\sum_{a\in B}S^{+}(u_a)+\sum_{a=1}^{l+m}S^{-}(v_a)}|i\rangle_B.
\end{aligned}\tag{4.22}$$

Using (3.15), we calculate the following vacuum expectation value :

$$\begin{aligned}
&{}_B\langle i|e^{\sum_{j=1}^{M}P(\zeta_j^2)-\sum_{j=1}^{N}P(q^{-1}\xi_j^2)+\sum_{a\in A}R^{-}(w_a)+\sum_{a\in B}R^{+}(u_a)+\sum_{a=1}^{l+m}R^{-}(v_a)}\\
\times\;&e^{\sum_{j=1}^{M}Q(\zeta_j^2)-\sum_{j=1}^{N}Q(q\xi_j^2)+\sum_{a\in A}S^{-}(w_a)+\sum_{a\in B}S^{+}(u_a)+\sum_{a=1}^{l+m}S^{-}(v_a)}|i\rangle_B\\
=\;&\langle i|e^{G^{(i)}}e^{\sum_{n=1}^{\infty}a_{-n}X_n}e^{-\sum_{n=1}^{\infty}a_nY_n}e^{F^{(i)}}|i\rangle.
\end{aligned}\tag{4.23}$$

Here we have defined

$$X_n=\frac{q^{7n/2}}{[2n]_q}\sum_{j=1}^{M}\zeta_j^{2n}-\frac{q^{5n/2}}{[2n]_q}\sum_{j=1}^{N}\xi_j^{2n}-\frac{q^{n/2}}{[n]_q}\sum_{a\in A}w_a^n+\frac{q^{-n/2}}{[n]_q}\sum_{a\in B}u_a^n-\frac{q^{n/2}}{[n]_q}\sum_{a=1}^{K}v_a^n, \tag{4.24}$$

$$Y_n = \frac{q^{-5n/2}}{[2n]_q}\sum_{j=1}^{M}\zeta_j^{-2n} - \frac{q^{-7n/2}}{[2n]_q}\sum_{j=1}^{N}\xi_j^{-2n} - \frac{q^{n/2}}{[n]_q}\sum_{a\in A}w_a^{-n} + \frac{q^{-n/2}}{[n]_q}\sum_{a\in B}u_a^{-n} - \frac{q^{n/2}}{[n]_q}\sum_{a=1}^{K}v_a^{-n}. \tag{4.25}$$

• The followings are the main results of this Section. For $K=|B|-|A|+\frac{1}{2}(M-N)<0$ we have

$$F^{(-,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)=0. \tag{4.26}$$

For $K=|B|-|A|+\frac{1}{2}(M-N)=0$, note that the form factors for triangular boundary conditions coincide with those for diagonal boundary conditions :

$$F^{(-,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)=F^{(i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r). \tag{4.27}$$

For $K=|B|-|A|+\frac{1}{2}(M-N)>0$, the following integral representations of the form factors of local operators is obtained:

$$\begin{aligned}
&F^{(-,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)\\
=\;&(s/q^2)^K(q^2/r)^{iK}q^{|B|}(1-q^2)^{|A|+2|B|}(-q^3)^{\frac{1}{4}(M-N)^2+\frac{i}{2}(M-N)-\sum_{a\in A}a-\sum_{b\in B}b}\\
\times\;&\left(\frac{\{q^6\}_\infty}{\{q^8\}_\infty}\right)^M\left(\frac{\{q^4\}_\infty}{\{q^6\}_\infty}\right)^N\{(q^4;q^2)_\infty\}^{2|B|+\frac{1}{2}(M-N)}\prod_{j=1}^{M}\zeta_j^{\frac{1+\epsilon_j}{2}+M-N-j+i}\prod_{j=1}^{N}\xi_j^{\frac{\mu_j+1}{2}+N-j+1-i}\\
\times\;&\prod_{1\le j<k\le M}\frac{\{q^6\zeta_j^2\zeta_k^2\}_\infty\{q^2/\zeta_j^2\zeta_k^2\}_\infty\{q^6\zeta_j^2/\zeta_k^2\}_\infty\{q^2\zeta_k^2/\zeta_j^2\}_\infty}{\{q^8\zeta_j^2\zeta_k^2\}_\infty\{q^4/\zeta_j^2\zeta_k^2\}_\infty\{q^8\zeta_j^2/\zeta_k^2\}_\infty\{q^4\zeta_k^2/\zeta_j^2\}_\infty}\\
\times\;&\prod_{1\le j<k\le N}\frac{\{q^4\xi_j^2\xi_k^2\}_\infty\{1/\xi_j^2\xi_k^2\}_\infty\{q^4\xi_j^2/\xi_k^2\}_\infty\{\xi_k^2/\xi_j^2\}_\infty}{\{q^6\xi_j^2\xi_k^2\}_\infty\{q^2/\xi_j^2\xi_k^2\}_\infty\{q^6\xi_j^2/\xi_k^2\}_\infty\{q^2\xi_k^2/\xi_j^2\}_\infty}\\
\times\;&\prod_{j=1}^{M}\prod_{k=1}^{N}\frac{\{q^7\zeta_j^2\xi_k^2\}_\infty\{q^3/\zeta_j^2\xi_k^2\}_\infty\{q^7\zeta_j^2/\xi_k^2\}_\infty\{q^3\xi_k^2/\zeta_j^2\}_\infty}{\{q^5\zeta_j^2\xi_k^2\}_\infty\{q/\zeta_j^2\xi_k^2\}_\infty\{q^5\zeta_j^2/\xi_k^2\}_\infty\{q\xi_k^2/\zeta_j^2\}_\infty}\\
\times\;&\prod_{j=1}^{M}\frac{[q^{10}\zeta_j^4]_\infty[q^{14}\zeta_j^4]_\infty[q^{10}/\zeta_j^4]_\infty[q^6/\zeta_j^4]_\infty}{[q^{12}\zeta_j^4]_\infty[q^{16}\zeta_j^4]_\infty[q^{12}/\zeta_j^4]_\infty[q^8/\zeta_j^4]_\infty}\prod_{j=1}^{N}\frac{[q^4\xi_j^4]_\infty[q^8\xi_j^4]_\infty[1/\xi_j^4]_\infty[q^4/\xi_j^4]_\infty}{[q^6\xi_j^4]_\infty[q^{10}\xi_j^4]_\infty[q^2/\xi_j^4]_\infty[q^6/\xi_j^4]_\infty}\\
\times\;&\sum_{\substack{l,m\ge 0\\ l+m=K}}\frac{(-1)^{m(M+1)}q^{\frac{l(l-1)-m(m-1)}{2}(2i-1)-3Mm}}{[l]_q![m]_q!}\oint\cdots\oint_{C_l^{(-,i)}}\prod_{a\in A}\frac{dw_a}{2\pi\sqrt{-1}}w_a^{1-i}\prod_{b\in B}\frac{du_b}{2\pi\sqrt{-1}}u_b^{i}\prod_{c=1}^{K}\frac{dv_c}{2\pi\sqrt{-1}}v_c^{1-2i}\\
\times\;&\prod_{a\in A}(w_a^2/q^2;q^4)_\infty(q^6/w_a^2;q^4)_\infty\prod_{a\in B}(u_a^2/q^2;q^4)_\infty(q^6/u_a^2;q^4)_\infty\prod_{a=1}^{K}(v_a^2/q^2;q^4)_\infty(q^6/v_a^2;q^4)_\infty\\
\times\;&\frac{\prod_{1\le a<b\le K}v_a^2(v_av_b/q^2;q^2)_\infty(q^4v_a/v_b;q^2)_\infty(v_b/v_a;q^2)_\infty(q^6/v_av_b;q^2)_\infty}{\prod_{a\in A}\prod_{b\in B}w_a^2(w_au_b/q^3;q^2)_\infty(q^3w_a/u_b;q^2)_\infty(u_b/qw_a;q^2)_\infty(q^5/w_au_b;q^2)_\infty}\\
\times\;&\frac{\prod_{a=1}^{K}\prod_{b\in A}(v_aw_b/q^2;q^2)_\infty(q^4v_a/w_b;q^2)_\infty(w_b/v_a;q^2)_\infty(q^6/v_aw_b;q^2)_\infty}{\prod_{a=1}^{K}\prod_{b\in B}v_a^2(v_au_b/q^2;q^2)_\infty(q^3v_a/u_b;q^2)_\infty(u_b/qv_a;q^2)_\infty(q^5/v_au_b;q^2)_\infty}\\
\times\;&\frac{\prod_{\substack{a,b\in A\\ a<b}}w_a^2(w_aw_b/q^2;q^2)_\infty(q^4w_a/w_b;q^2)_\infty(w_b/w_a;q^2)_\infty(q^6/w_aw_b;q^2)_\infty}{\prod_{a\in A}\left\{\prod_{1\le j\le a}(\zeta_j^2-q^{-2}w_a)\prod_{a\le j\le M}(w_a-q^4\zeta_j^2)\right\}}\\
\times\;&\frac{\prod_{\substack{a,b\in B\\ a<b}}u_a^2(u_au_b/q^4;q^2)_\infty(q^2u_a/u_b;q^2)_\infty(u_b/q^2u_a;q^2)_\infty(q^4/u_au_b;q^2)_\infty}{\prod_{a\in B}\left\{\prod_{1\le j\le a}(\xi_j^2-q^{-4}u_a)\prod_{a\le j\le M}(u_a-q^2\xi_j^2)\right\}}
\end{aligned}$$

$$
\begin{aligned}
&\times \frac{\prod_{j=1}^{M}\prod_{a\in B}u_a(q\zeta_j^2u_a;q^4)_\infty(q^3\zeta_j^2/u_a;q^4)_\infty(qu_a/\zeta_j^2;q^4)_\infty(q^3/\zeta_j^2u_a;q^4)_\infty}{\prod_{j=1}^{M}\prod_{a\in A}(q^2\zeta_j^2w_a;q^4)_\infty(q^8\zeta_j^2/w_a;q^4)_\infty(q^2w_a/\zeta_j^2;q^4)_\infty(q^4/\zeta_j^2w_a;q^4)_\infty}\\
&\times \frac{\prod_{j=1}^{N}\prod_{a=1}^{K}v_a(q\xi_j^2v_a;q^4)_\infty(q^3\xi_j^2/v_a;q^4)_\infty(qv_a/\xi_j^2;q^4)_\infty(q^3/\xi_j^2v_a;q^4)_\infty}{\prod_{j=1}^{M}\prod_{a=1}^{K}(q^2\zeta_j^2v_a;q^4)_\infty(q^8\xi_j^2/v_a;q^4)_\infty(q^2v_a/\xi_j^2;q^4)_\infty(q^4/\xi_j^2v_a;q^4)_\infty}\\
&\times \frac{\prod_{a\in A}\left\{\prod_{b=1}^{l}(v_b-w_a)(v_b-q^2w_a)\prod_{b=l+1}^{K}(w_a-v_b)(w_a-q^2v_b)\right\}}{\prod_{a=1}^{M}\left\{\prod_{a=1}^{l}(v_a-q^4\zeta_j^2)\prod_{a=l+1}^{K}(\zeta_j^2-q^{-2}v_a)\right\}}\\
&\times \begin{cases}\prod_{j=1}^{M}\frac{(q^2r\zeta_j^2;q^4)_\infty}{(q^4r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q^3r\xi_j^2;q^4)_\infty}{(qr\xi_j^2;q^4)_\infty}\frac{\prod_{a\in B}(1-ru_a/q^3)}{\prod_{a\in A}(1-rw_a/q^2)\prod_{a=1}^{K}(1-rv_a/q^2)} & (i=0),\\ \prod_{j=1}^{M}\frac{(1/r\zeta_j^2;q^4)_\infty}{(q^2/r\zeta_j^2;q^4)_\infty}\prod_{j=1}^{N}\frac{(q/r\xi_j^2;q^4)_\infty}{(1/qr\xi_j^2;q^4)_\infty}\frac{\prod_{a\in B}(1-q/ru_a)}{\prod_{a\in A}(1-q^2/rw_a)\prod_{a=1}^{K}(1-q^2/rv_a)} & (i=1).\end{cases}
\end{aligned}
\tag{4.28}
$$

Here the integration contour $C_l^{(-,0)}$ is a closed curve that satisfies the following conditions for $s=0,1,2,\cdots$. The w_a $(a\in A)$ encircles $q^{8+4s}\zeta_j^2$ $(1\le j<a)$, $q^{4+4s}\zeta_j^2$ $(a\le j\le M)$, q^{4+4s}/ζ_j^2 $(1\le j\le M)$, $q^{-1+2s}u_b$, q^{5+2s}/u_b $(b\in B)$, but not $q^{2-4s}\zeta_j^2$ $(1\le j\le a)$, $q^{-2-4s}\zeta_j^2$ $(a<j\le M)$, q^{-2-4s}/ζ_j^2 $(1\le j\le M)$, $q^{-3-2s}u_b$, q^{3-2s}/u_b $(b\in B)$, q^2/r. The u_b $(b\in B)$ encircles $q^{6+4s}\xi_j^2$ $(1\le j<b)$, $q^{2+4s}\xi_j^2$ $(b\le j\le N)$, q^{2+4s}/ξ_j^2 $(1\le j\le N)$, $q^{3+2s}w_a$, q^{5+2s}/w_a $(a\in B)$, $q^{3+2s}v_c$ $(1\le c\le l)$, $q^{-1+2s}v_c$ $(l<c\le K)$, q^{5+2s}/v_c $(1\le c\le K)$, but not $q^{4-4s}\xi_j^2$ $(1\le j\le b)$, $q^{-4s}\xi_j^2$ $(b<j\le N)$, q^{-4s}/ξ_j^2 $(1\le j\le N)$, $q^{1-2s}w_a$, q^{-3-2s}/w_a $(a\in A)$, $q^{1-2s}v_c$ $(1\le c\le l)$, $q^{-3-2s}v_c$ $(l<c\le K)$, q^{3-2s}/v_c $(1\le c\le K)$. The v_c $(1\le c\le l)$ encircles $q^{4+4s}\zeta_j^2$, q^{4+4s}/ζ_j^2 $(1\le j\le M)$, $q^{-1+2s}u_b$, $q^{5+2s}u_b$ $(b\in B)$, but not $q^{-2-4s}\zeta_j^2$, q^{-2-4s}/ζ_j^2 $(1\le j\le M)$, $q^{-3-2s}u_b$, $q^{3-2s}u_b$ $(b\in B)$, q^2/r. The v_c $(l<c\le K)$ encircles $q^{8+4s}\zeta_j^2$, q^{4+4s}/ζ_j^2 $(1\le j\le M)$, $q^{3+2s}u_b$, $q^{5+2s}u_b$ $(b\in B)$, but not $q^{2-4s}\zeta_j^2$, q^{-2-4s}/ζ_j^2 $(1\le j\le M)$, $q^{1-2s}u_b$, $q^{3-2s}u_b$ $(b\in B)$, q^2/r. Here the integration contour $C_l^{(-,0)}$ is a closed curve such that w_a $(a\in A)$ encircles q^2/r in addition the same points $C_l^{(-,0)}$ does.

4.4 Identities between multiple integrals

From the spin-reversal properties in (2.37), (2.39), and (2.40), we have the following relations between form factors of local operators :

$$
\begin{aligned}
&F^{(\pm,i,M,N)}_{\epsilon_1,\cdots,\epsilon_M,\mu_1,\cdots,\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;r,s)\\
=\;& F^{(\mp,1-i,M,N)}_{-\epsilon_1,\cdots,-\epsilon_M,-\mu_1,\cdots,-\mu_N}(\zeta_1,\cdots,\zeta_M,\xi_1,\cdots,\xi_N;1/r,-s/r).
\end{aligned}
\tag{4.29}
$$

We should understand this spin-reversal property (4.29) as an analytic continuation of the parameter r. From the above (4.29), similarly to the diagonal case we obtain infinitely many relations between n-fold integrals which cannot be reduced to the relations between n-fold integrals of elliptic gamma functions summarized in [41]. Here we focus on a simple example.

• From the following relation between form factors :

$$
F^{(+,1,2,0)}_{+,+}(-q^{-1}\zeta,\zeta;r,s)=F^{(-,0,2,0)}_{-,-}(-q^{-1}\zeta,\zeta;1/r,-s/r),
\tag{4.30}
$$

we deduce the identity:

$$
2+\frac{1-\zeta^2/r}{\zeta^2}\sum_{k=1}^{\infty}(-q^2)^k\frac{(\zeta^2-\zeta^{-2})-(1+q^{4k})/r+(\zeta^2+\zeta^{-2})q^{2k}}{(1-q^{2k}\zeta^2/r)(1-q^{2k}/r\zeta^2)}
$$

$$
\begin{aligned}
&= q^2\frac{(q^2;q^2)_\infty^8}{(q^4;q^4)_\infty^4}\frac{\Theta_{q^4}(\zeta^4)}{1-\zeta^4}\left(q^2\oint\oint\oint_{C_0^{(+,1)}}-\oint\oint\oint_{C_1^{(+,1)}}\right)\prod_{a=1}^{3}\frac{dw_a}{2\pi\sqrt{-1}}\\
&\times\ \frac{(1-1/r\zeta^2)(1-q/rw_3)\prod_{a=1}^{2}(1-q^2/\zeta^2w_a)}{w_2^2w_3^3(1-q^2w_1/w_2)(1-q^4/w_1w_2)\prod_{a=1,2}(1-q^2/rw_a)}\\
&\times\ \frac{\Theta_{q^2}(w_1w_2)\Theta_{q^2}(w_2/w_1)\Theta_{q^2}(\zeta^2w_3/q)\Theta_{q^2}(qw_3/\zeta^2)\prod_{a=1}^{3}\Theta_{q^4}(w_a^2/q^2)}{\prod_{a=1}^{2}\Theta_{q^2}(w_aw_3/q^2)\Theta_{q^2}(w_a/qw_3)\Theta_{q^2}(w_a\zeta^2)\Theta_{q^2}(w_a/\zeta^2)}.
\end{aligned}
\tag{4.31}
$$

The integration contour $C_0^{(+,1)}$ is a simple closed curve such that the w_1 encircles $q^{2+2s}\zeta^2$, q^{4+2s}/ζ^2, $q^{-1+2s}w_3$, q^{5+2s}/w_3, q^2/r the w_2 encircles $q^{4+2s}\zeta^2$, q^{4+2s}/ζ^2, $q^{-1+2s}w_3$, q^{5+2s}/w_3, q^2/r, and the w_3 encircles $q^{3+2s}w_a, q^{5+2s}/w_a$ $(a=1,2)$, for $s=0,1,2,\cdots$. The integration contour $C_1^{(+,1)}$ is a simple closed curve such that the w_1 encircles $q^{2+2s}\zeta^2$, q^{4+2s}/ζ^2, $q^{3+2s}w_3$, q^{5+2s}/w_3, the w_2 encircles $q^{4+2s}\zeta^2$, q^{4+2s}/ζ^2, $q^{3+2s}w_3$, q^{5+2s}/w_3, and w_3 encircles $q^{-1+2s}w_a, q^{5+2s}/w_a$ $(a=1,2)$, for $s=0,1,2,\cdots$.

5 Concluding remarks

In the present article, integral representations of form factors of local operators of the half-infinite XXZ spin chain for diagonal or triangular boundary conditions in the massive regime have been proposed. Completing the analysis of [23, 21, 24], they are written in the form of multiple integrals of meromorphic functions. For some simple relations between the number of type I and type II vertex operators inserted, the expressions for diagonal or triangular boundary conditions are identical. For diagonal boundary conditions, the expressions here obtained essentially coincide with those derived from the QISM [18, 19]. In addition, the constructive approach introduced in [24] which allows to derive identities between n-fold multiple integrals of elliptic theta functions (obtained from expectation values of type I vertex operators) has been extended to the family of multiple integrals associated with form factors: corresponding identities follow from expectation values of combinations of type I and type II vertex operators. A direct proof of such remarkable identities would be highly desirable.

For more general integrable boundary conditions, the explicit construction of the vacuum eigenvectors and dual ones starting from the representation theory of the q-Onsager algebra can be considered following [21]. This is currently investigated. As soon as the eigenvectors and duals will be identified, the VOA to the computation of correlation functions and form factors can be applied in a straightforward manner, extending the results of [23, 24] and the ones here presented.

For models with higher symmetries [6, 7, 9, 10, 11, 12], recall that the VOA approach has been applied successfully for diagonal boundary conditions in some cases [33, 34, 35]. For the case of models associated with higher symmetries and more general boundary conditions, the analysis presented in [21, 24] and here may be considered. To this end, let us mention that the non-Abelian infinite dimensional algebra which occurs in the thermodynamic limit is a coideal subalgebra of the basic quantum group. For the family of affine Lie algebras $\widehat{g}$, coideal subalgebras of $U_q(\widehat{g})$ generate the so-called generalized q-Onsager algebras introduced in [36]. As a starting point, infinite dimensional (q-vertex operators) representations have to be considered, following [22, 21].

To conclude, we would like to stress that alternative derivations of the expressions here proposed would

be highly desirable. Based on the recent results for a triangular boundary [14], this may be feasible using the QISM. For more general integrable boundary conditions, other approaches than the ones based on Onsager's formulation/VOA would be required: for instance, a modified algebraic BA approach for the XXZ open spin chain inspired by [37, 38], the Sklyanin's separation of variables approach [40] or the approach developed in [39] seem to be promising.

Acknowledgements

T.K. would like to thank Laboratoire de Mathématiques et Physique Théorique, Université de Tours for kind invitation and warm hospitality during his stay in March 2013. This work is supported by the Grant-in-Aid for Scientific Research **C** (21540228) from JSPS and Visiting professorship from CNRS.

A Definitions of the R and K matrices

In Sklyanin's framework [13], the transfer matrix associated with a finite spin chain is built from two objects: the R-matrix and the $K-$matrix. For the model (2.2), the R-matrix $R(\zeta)$ defined by:

$$R(\zeta)=\frac{1}{\kappa(\zeta)}\begin{pmatrix}1&&&\\&\frac{(1-\zeta^2)q}{1-q^2\zeta^2}&\frac{(1-q^2)\zeta}{1-q^2\zeta^2}&\\&\frac{(1-q^2)\zeta}{1-q^2\zeta^2}&\frac{(1-\zeta^2)q}{1-q^2\zeta^2}&\\&&&1\end{pmatrix}, \tag{A.1}$$

where we have set

$$\kappa(\zeta)=\zeta\frac{(q^4\zeta^2;q^4)_\infty(q^2/\zeta^2;q^4)_\infty}{(q^4/\zeta^2;q^4)_\infty(q^2\zeta^2;q^4)_\infty},\quad (z;p)_\infty=\prod_{n=0}^{\infty}(1-p^nz). \tag{A.2}$$

As an operator on $V\otimes V$, the matrix elements of $R(\zeta)\in\mathrm{End}(V\otimes V)$ are given by $R(\zeta)v_{\epsilon_1}\otimes v_{\epsilon_2}=\sum_{\epsilon_1',\epsilon_2'=\pm}v_{\epsilon_1'}\otimes v_{\epsilon_2'}R(\zeta)_{\epsilon_1'\epsilon_2'}^{\epsilon_1\epsilon_2}$, where the ordering of the index is given by $v_+\otimes v_+, v_+\otimes v_-, v_-\otimes v_+, v_-\otimes v_-$. As usual, when copies V_j of V are involved, $R_{ij}(\zeta)$ acts as $R(\zeta)$ on the i-th and j-th components and as identity elsewhere. By definition, the R-matrix $R(\zeta)$ satisfies the Yang-Baxter equation :

$$R_{12}(\zeta_1/\zeta_2)R_{13}(\zeta_1/\zeta_3)R_{23}(\zeta_2/\zeta_3)=R_{23}(\zeta_2/\zeta_3)R_{13}(\zeta_1/\zeta_3)R_{12}(\zeta_1/\zeta_2). \tag{A.3}$$

The normalization factor $\kappa(\zeta)$ (A.2) is determined by the following unitarity and crossing symmetry conditions:

$$R_{12}(\zeta)R_{21}(\zeta^{-1})=1,\quad R(\zeta)_{\epsilon_2\epsilon_1'}^{\epsilon_2'\epsilon_1}=R(-q^{-1}\zeta^{-1})_{-\epsilon_1\epsilon_2}^{-\epsilon_1'\epsilon_2'}. \tag{A.4}$$

The second basic object is the triangular K-matrix: $K^{(\pm)}(\zeta)=K^{(\pm)}(\zeta;r,s)$ [28, 29] by

$$K^{(+)}(\zeta;r,s) = \frac{\varphi(\zeta^2;r)}{\varphi(\zeta^{-2};r)}\begin{pmatrix}\frac{1-r\zeta^2}{\zeta^2-r}&\frac{s\zeta(\zeta^2-\zeta^{-2})}{\zeta^2-r}\\0&1\end{pmatrix}, \tag{A.5}$$

$$K^{(-)}(\zeta;r,s) = \frac{\varphi(\zeta^2;r)}{\varphi(\zeta^{-2};r)}\begin{pmatrix}\frac{1-r\zeta^2}{\zeta^2-r}&0\\\frac{s\zeta(\zeta^2-\zeta^{-2})}{\zeta^2-r}&1\end{pmatrix}, \tag{A.6}$$

where we have set

$$\varphi(z;r)=\frac{(q^4rz;q^4)_\infty(q^6z^2;q^8)_\infty}{(q^2rz;q^4)_\infty(q^8z^2;q^8)_\infty}. \tag{A.7}$$

When viewed as an operator on V, the matrix elements of $K^{(\pm)}(\zeta)\in \mathrm{End}(V)$ are given by $K^{(\pm)}(\zeta)v_\epsilon=\sum_{\epsilon'=\pm}v_{\epsilon'}K^{(\pm)}(\zeta)^{\epsilon}_{\epsilon'}$, where the ordering of the index is given by v_+,v_-. As usual, when copies V_j of V are involved, $K^{(\pm)}_j(\zeta)$ acts as $K^{(\pm)}(\zeta)$ on the j-th component and as identity elsewhere. The K-matrix $K^{(\pm)}(\zeta)$ satisfies the boundary Yang-Baxter equation (also called the reflection equation):

$$K^{(\pm)}_2(\zeta_2)R_{21}(\zeta_1\zeta_2)K^{(\pm)}_1(\zeta_1)R_{12}(\zeta_1/\zeta_2)=R_{21}(\zeta_1/\zeta_2)K^{(\pm)}_1(\zeta_1)R_{12}(\zeta_1\zeta_2)K^{(\pm)}_2(\zeta_2). \tag{A.8}$$

The normalization factor (2.7) is determined by the following boundary unitarity and boundary crossing symmetry [29]:

$$K^{(\pm)}(\zeta)K^{(\pm)}(\zeta^{-1})=1,\quad K^{(\pm)}(-q^{-1}\zeta^{-1})^{\epsilon_2}_{\epsilon_1}=\sum_{\epsilon_1',\epsilon_2'=\pm}R(-q\zeta^2)^{-\epsilon_1\epsilon_2}_{\epsilon_1'-\epsilon_2'}K^{(\pm)}(\zeta)^{\epsilon_1'}_{\epsilon_2'}. \tag{A.9}$$

The $K^{(\pm)}(\zeta;r,s)$ defined in (A.5) and (A.6) give general scalar triangular solutions of (A.8) and (A.9). For $s=0$, one recovers the scalar diagonal solution of the reflection equation.

B Bosonizations of vertex operators

In this Appendix we review the bosonizations of the quantum group $U_q(\widehat{sl_2})$ and the q-vertex operators for level $k=1$ [5, 8]. The center $\gamma^{\frac{1}{2}}$ of $U_q(\widehat{sl_2})$ satisfies $(\gamma^{\frac{1}{2}})^2=q^{h_0+h_1}=q^k$ on the level k representation. Hence $\gamma=q$ on the level $k=1$ representation. For the level $k=1$ case, the defining relation of a_m ($m\in\mathbf{Z}_{\neq 0}$) becomes

$$[a_m,a_n]=\delta_{m+n,0}\frac{[2m]_q[m]_q}{m}\quad (m,n\neq 0). \tag{B.1}$$

We introduce the zero-mode operator ∂,α by $[\partial,\alpha]=2$ and the normal ordering symbol $:\ :$

$$:a_ma_n:=\begin{cases}a_ma_n & (m<0)\\ a_na_m & (m>0)\end{cases},\qquad :\alpha\partial:=:\partial\alpha:=\alpha\partial. \tag{B.2}$$

The bosonizations of the irreducible highest representation $V(\Lambda_i)$ and its dual $V^*(\Lambda_i)$ with fundamental weight Λ_i are given by

$$V(\Lambda_i) = \mathbf{C}[a_{-1},a_{-2},a_{-3}\cdots]\otimes\left(\oplus_{n\in\mathbf{Z}}\mathbf{C}e^{\Lambda_i+n\alpha}\right), \tag{B.3}$$

$$V^*(\Lambda_i) = \mathbf{C}[a_1,a_2,a_3,\cdots]\otimes\left(\oplus_{n\in\mathbf{Z}}\mathbf{C}e^{-\Lambda_i-n\alpha}\right). \tag{B.4}$$

On this space the actions of a_m, e^α, ∂ are given by

$$a_m(f\otimes e^\beta)=\begin{cases}a_mf\otimes e^\beta & (m<0),\\ [a_m,f]\otimes e^\beta & (m>0),\end{cases} \tag{B.5}$$

$$e^\alpha(f\otimes e^\beta)=f\otimes e^{\alpha+\beta},\quad \partial(f\otimes e^\beta)=(\alpha,\beta)f\otimes e^\beta, \tag{B.6}$$

where $f \in \mathbf{C}[a_{-1}, a_{-2}, \cdots]$, and we have set $[\partial, \Lambda_0] = 0$ and $\Lambda_1 = \Lambda_0 + \frac{\alpha}{2}$.

The action of other Drinfeld generators is given by

$$K = q^{\partial}, \quad \gamma = q, \tag{B.7}$$

$$X^{\pm}(z) = \exp\left(R^{\pm}(z)\right) \exp\left(S^{\pm}(z)\right) e^{\pm\alpha} z^{\pm\partial}, \tag{B.8}$$

$$q^d(1 \otimes e^{\Lambda_i + n\alpha}) = q^{-(\beta,\beta)/2 + i/4}(1 \otimes e^{\Lambda_i + n\alpha}), \tag{B.9}$$

where

$$R^{\pm}(z) = \pm \sum_{n=1}^{\infty} \frac{a_{-n}}{[n]_q} q^{\mp n/2} z^n, \quad S^{\pm}(z) = \mp \sum_{n=1}^{\infty} \frac{a_n}{[n]_q} q^{\mp n/2} z^{-n}. \tag{B.10}$$

We have the following bosonizations of the q-vertex operators :

$$\Phi_{-}^{(1-i,i)}(\zeta) = \exp\left(P(\zeta^2)\right) \exp\left(Q(\zeta^2)\right) e^{\alpha/2} (-q^3\zeta^2)^{(\partial+i)/2} \zeta^{-i}, \tag{B.11}$$

$$\Phi_{+}^{(1-i,i)}(\zeta) = \oint_{\widetilde{C}_1} \frac{dw}{2\pi\sqrt{-1}} \frac{(1-q^2)w\zeta}{q(w-q^2\zeta^2)(w-q^4\zeta^2)} : \Phi_{-}^{(1-i,i)}(\zeta) X^{-}(w) :, , \tag{B.12}$$

$$\Psi_{-}^{*(1-i,i)}(\xi) = \exp\left(-P(q^{-1}\xi^2)\right) \exp\left(-Q(q\xi^2)\right) e^{-\alpha/2} (-q^3\xi^2)^{(-\partial+i)/2} \xi^{1-i}, \tag{B.13}$$

$$\Psi_{+}^{*(1-i,i)}(\xi) = \oint_{\widetilde{C}_2} \frac{dw}{2\pi\sqrt{-1}} \frac{q^2(1-q^2)\xi}{(w-q^2\xi^2)(w-q^4\xi^2)} : \Psi_{-}^{*(1-i,i)}(\xi) X^{+}(w) :, \tag{B.14}$$

where

$$P(z) = \sum_{n=1}^{\infty} \frac{a_{-n}}{[2n]_q} q^{7n/2} z^n, \quad Q(z) = -\sum_{n=1}^{\infty} \frac{a_n}{[2n]_q} q^{-5n/2} z^{-n}. \tag{B.15}$$

Here the integration contour $\widetilde{C}_1$ is a simple closed curve such that the w encircles $q^4\zeta^2$ inside but not $q^2\zeta^2$. The integration contour $\widetilde{C}_2$ is a simple closed curve such that the w encircles $q^2\zeta^2$ inside but not $q^4\zeta^2$.

In what follows we summarize normal orderings.

$$\Phi_{-}^{(i,1-i)}(\zeta_1)\Phi_{-}^{(1-i,i)}(\zeta_2) = (-q^3\zeta_1^2)^{\frac{1}{2}} \frac{(q^2\zeta_2^2/\zeta_1^2; q^4)_\infty}{(q^4\zeta_2^2/\zeta_1^2; q^4)_\infty} : \Phi_{-}^{(i,1-i)}(\zeta_1)\Phi_{-}^{(1-i,i)}(\zeta_2) :, \tag{B.16}$$

$$\Psi_{-}^{*(i,1-i)}(\xi_1)\Psi_{-}^{*(1-i,i)}(\xi_2) = (-q^3\xi_1^2)^{\frac{1}{2}} \frac{(\xi_2^2/\xi_1^2; q^4)_\infty}{(q^2\xi_2^2/\xi_1^2; q^4)_\infty} : \Psi_{-}^{*(1-i,i)}(\xi_1)\Psi_{-}^{*(1-i,i)}(\xi_2) :, \tag{B.17}$$

$$\Phi_{-}^{(i,1-i)}(\zeta)\Psi_{-}^{*(1-i,i)}(\xi) = (-q^3\zeta^2)^{-\frac{1}{2}} \frac{(q^3\xi^2/\zeta^2; q^4)_\infty}{(q\xi^2/\zeta^2; q^4)_\infty} : \Phi_{-}^{(i,1-i)}(\zeta)\Psi_{-}^{*(1-i,i)}(\xi) :, \tag{B.18}$$

$$\Psi_{-}^{*(i,1-i)}(\xi)\Phi_{-}^{(1-i,i)}(\zeta) = (-q^3\xi^2)^{-\frac{1}{2}} \frac{(q^3\zeta^2/\xi^2; q^4)_\infty}{(q\zeta^2/\xi^2; q^4)_\infty} : \Psi_{-}^{*(i,1-i)}(\xi)\Phi_{-}^{(1-i,i)}(\zeta) :, \tag{B.19}$$

$$X^{+}(w_1)X^{+}(w_2) = w_1^2(1 - w_2/w_1)(1 - w_2/q^2w_1) : X^{+}(w_1)X^{+}(w_2) :, \tag{B.20}$$

$$X^{-}(w_1)X^{-}(w_2) = w_1^2(1 - w_2/w_1)(1 - q^2w_2/w_1) : X^{-}(w_1)X^{-}(w_2) :, \tag{B.21}$$

$$X^{+}(w_1)X^{-}(w_2) = \frac{1}{w_1^2(1 - qw_2/w_1)(1 - w_2/qw_1)} : X^{+}(w_1)X^{-}(w_2) :, \tag{B.22}$$

$$X^{-}(w_1)X^{+}(w_2) = \frac{1}{w_1^2(1 - qw_2/w_1)(1 - w_2/qw_1)} : X^{-}(w_1)X^{+}(w_2) :, \tag{B.23}$$

$$\Phi_{-}^{(i,1-i)}(\zeta)X^{+}(w) = (-q^3\zeta^2)(1 - w/q^3\zeta^2) : \Phi_{-}^{(i,1-i)}(\zeta)X^{+}(w) :, \tag{B.24}$$

$$X^+(w)\Phi_-^{(i,1-i)}(\zeta) = w(1-q^3\zeta^2/w) : X^+(w)\Phi_-^{(i,1-i)}(\zeta) :, \tag{B.25}$$

$$\Phi_-^{(i,1-i)}(\zeta)X^-(w) = \frac{-1}{q^3\zeta^2(1-w/q^2\zeta^2)} : \Phi_-^{(i,1-i)}(\zeta)X^-(w) :, \tag{B.26}$$

$$X^-(w)\Phi_-^{(i,1-i)}(\zeta) = \frac{1}{w(1-q^4\zeta^2/w)} : X^-(w)\Phi_-^{(i,1-i)}(\zeta) :, \tag{B.27}$$

$$\Psi_-^{*(i,1-i)}(\xi)X^+(w) = \frac{-1}{q^3\xi^2(1-w/q^4\xi^2)} : \Psi_-^{*(i,1-i)}(\xi)X^+(w) :, \tag{B.28}$$

$$X^+(w)\Psi_-^{*(i,1-i)}(\xi) = \frac{1}{w(1-q^2\xi^2/w)} : X^+(w)\Psi_-^{*(i,1-i)}(\xi) :, \tag{B.29}$$

$$\Psi_-^{*(i,1-i)}(\xi)X^-(w) = (-q^3\xi^2)(1-w/q^3\xi^2) : \Psi_-^{*(i,1-i)}(\xi)X^-(w) :, \tag{B.30}$$

$$X^-(w)\Psi_-^{*(i,1-i)}(\xi) = w(1-q^3\xi^2/w) : X^-(w)\Psi_-^{*(i,1-i)}(\xi) : . \tag{B.31}$$

C Proof of the spin-reversal property

In this Appendix we give a direct proof of one of the simplest example of the spin-reversal property. We would like to show

$$F_{-,+}^{(1,2,0)}(-q^{-1}\zeta,\zeta;r) = F_{+,-}^{(0,2,0)}(-q^{-1}\zeta,\zeta;1/r). \tag{C.1}$$

We have

$$F_{+,-}^{(i,2,0)}(-q^{-1}\zeta,\zeta;r) = \frac{(q^2;q^2)_\infty^3\Theta_{q^4}(\zeta^4)}{(1-\zeta^4)}\oint_{C_{+-}^{(i)}}\frac{dw}{2\pi i}\frac{(1-r\zeta^2)(1-\zeta^2w/q^2)}{\zeta^2(1-rw/q^2)}\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(\zeta^2w)\Theta_{q^2}(w/\zeta^2)}. \tag{C.2}$$

Here $C_{+-}^{(0)}$ is a closed curve satisfying that the w encircles $q^{2+2s}\zeta^2,\ q^{2+2s}/\zeta^2\ (s=0,1,2,\cdots)$ but not $q^{-2s}\zeta^2,\ q^{-2s}/\zeta^2\ (s=0,1,2,\cdots),\ q^2/r$. Here $C_{+-}^{(1)}$ is a closed curve such that w encircles q^2/r in addition the same points $C_{+-}^{(0)}$ does. Upon the specialization we have

$$F_{+,-}^{(i,2,0)}(-q^{-1}\zeta,\zeta;r) = (-)\frac{(q^2;q^2)_\infty^3\Theta_{q^4}(z^2)}{(1-\zeta^4)}\oint_{C_{-+}^{(i)}}\frac{dw}{2\pi i}\frac{(1-r\zeta^2)(1-\zeta^2w/q^2)}{\zeta^2(1-rw/q^2)}\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(\zeta^2w)\Theta_{q^2}(w/\zeta^2)}. \tag{C.3}$$

Here $C_{-+}^{(0)}$ is closed curve satisfying that the w encircles $q^{4+2s}\zeta^2,\ q^{4+2s}/\zeta^2\ (s=0,1,2,\cdots)$ but not $q^{2-2s}\zeta^2,\ q^{2-2s}/\zeta^2\ (s=0,1,2,\cdots),\ q^2/r$. Here $C_{-+}^{(1)}$ is closed curve satisfying that the w encircles q^2/r in addition the same points $C_{-+}^{(0)}$ does. We note that the integrand does not have a pole at $w=q^2/\zeta^2$. We have already deformed $C_{-+}^{(i)}$ from the original definition of the correlation functions at $w=q^2/\zeta^2$. Let's study

$$F_{+,-}^{(0,2,0)}(-q^{-1}\zeta,\zeta;r) - F_{-,+}^{(1,2,0)}(-q^{-1}\zeta,\zeta;1/r). \tag{C.4}$$

Changing $w\to q^4/w$ in the second term, we have

$$\frac{(q^2;q^2)_\infty^3\Theta_{q^4}(\zeta^4)}{\zeta^2}\oint_{C_{+-}^{(0)}}\frac{dw}{2\pi i}\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(w\zeta^2)\Theta_{q^2}(w/\zeta^2)}. \tag{C.5}$$

Taking the residues, we have

$$\oint_{C_{+-}^{(0)}}\frac{dw}{2\pi i}\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(w\zeta^2)\Theta_{q^2}(w/\zeta^2)} = \sum_{n=1}^{\infty}(-q^2)^n\left(\mathrm{Res}_{w=\zeta^2}+\mathrm{Res}_{w=1/\zeta^2}\right)\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(w\zeta^2)\Theta_{q^2}(w/\zeta^2)} = 0. \tag{C.6}$$

Here we have used $\frac{\Theta_{q^4}(q^{2-4n}w^2)}{\Theta_{q^2}(q^{-2n}w\zeta^2)\Theta_{q^2}(q^{-2n}w/\zeta^2)} = (-q^2)^n\frac{\Theta_{q^4}(q^2w^2)}{\Theta_{q^2}(w\zeta^2)\Theta_{q^2}(w/\zeta^2)}$.

D Convenient formulae

In this Appendix we summarize convenient formulae for calculations of vacuum expectation values.

$$\exp\left(\sum_{n=1}^{\infty}\frac{1}{(q-q^{-1})}\frac{q^{(c+2)n}}{n[2n]_q}y^n\right) = (q^{c+4}y;q^4)_\infty, \tag{D.1}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{1}{(q-q^{-1})}\frac{[n]_q q^{(c+2)n}}{n[2n]_q^2}y^n\right) = \frac{(q^{c+5}y;q^4,q^4)_\infty}{(q^{c+7}y;q^4,q^4)_\infty}, \tag{D.2}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{1}{(q-q^{-1})}\frac{q^{(c+2)n}}{n[n]_q}y^n\right) = (q^{c+3}y;q^4)_\infty, \tag{D.3}$$

$$\exp\left(\sum_{n>0}\frac{[n]_q}{n[2n]_q}\frac{q^{cn}}{1-\alpha_n\gamma_n}y^n\right) = \frac{(q^{c+3}y;q^4,q^4)_\infty}{(q^{c+1}y;q^4,q^4)_\infty}, \tag{D.4}$$

$$\exp\left(\sum_{n>0}\frac{[2n]_q}{n[n]_q}\frac{q^{cn}}{1-\alpha_n\gamma_n}y^n\right) = \frac{1}{(q^{c-1}y;q^2)_\infty}. \tag{D.5}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{[n]_q}{n(1-\alpha_n\gamma_n)}\delta_n^{(i)}q^{cn/2}y^n\right) = \left(\frac{(q^{c+3}y^2;q^8,q^8)_\infty(q^{c+1}y^2;q^8,q^8)_\infty}{(q^{c-1}y^2;q^8,q^8)_\infty(q^{c+5}y^2;q^8,q^8)_\infty}\right)^{1/2} \times \left(\frac{(q^{(c-3)/2+2i}r^{1-2i}y;q^4,q^4)_\infty}{(q^{(c+1)/2+2i}r^{1-2i}y;q^4,q^4)_\infty}\right)^{1-2i}, \tag{D.6}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{[2n]_q}{n(1-\alpha_n\gamma_n)}\delta_n^{(i)}q^{cn/2}y^n\right) = \left(\frac{(q^{c-1}y^2;q^8)_\infty}{(q^{c-3}y^2;q^8)_\infty}\right)^{1/2}(q^{(c-5)/2+2i}r^{1-2i}y;q^4)_\infty^{1-2i}, \tag{D.7}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{[n]_q}{n(1-\alpha_n\gamma_n)}\beta_n^{(i)}q^{cn/2}y^n\right) = \left(\frac{(q^{c+7}y^2;q^8,q^8)_\infty(q^{c+13}y^2;q^8,q^8)_\infty}{(q^{c+9}y^2;q^8,q^8)_\infty(q^{c+11}y^2;q^8,q^8)_\infty}\right)^{1/2} \times \left(\frac{(q^{(c+9)/2-2i}r^{1-2i}y;q^4,q^4)_\infty}{(q^{(c+13)/2-2i}r^{1-2i}y;q^4,q^4)_\infty}\right)^{1-2i}, \tag{D.8}$$

$$\exp\left(\sum_{n=1}^{\infty}\frac{[2n]_q}{n(1-\alpha_n\gamma_n)}\beta_n^{(i)}q^{cn/2}y^n\right) = \left(\frac{(q^{c+5}y^2;q^8)_\infty}{(q^{c+7}y^2;q^8)_\infty}\right)^{1/2}(q^{(c+7)/2-2i}r^{1-2i}y;q^4)_\infty^{1-2i}. \tag{D.9}$$

Here we have used α_n, γ_n, $\beta_n^{(i)}$, and $\delta_n^{(i)}$ in (2.25), (2.26), and (2.27).

References

[1] H.A.Bethe, Zur theorie der metalle I. eigenfunktionen der linearen atomkette, *Zeitschrift für Physik* **71**, 205-26 (1931)

[2] R.J.Baxter, *Exactly solved models in statistical mechanics*, Academic Press, London, 1982.

[3] V.E.Korepin, N.M.Bogoliubov, and A.G.Izergin, *Quantum inverse scattering method and correlation functions*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1993.

A.G.Izergin and V.E.Korepin, The quantum inverse scattering method approach to correlation functions, *Comm.Math.Phys.* **94**, 67-92, (1984).

A.G.Izergin and V.E.Korepin, Correlation functions for the Heisenberg XXZ-antiferromagnet, *Comm.Math.Phys.* **99**, 271-302, (1985).

[4] N.Kitanine, J.M.Maillet, and V.Terras, Form factors of the XXZ Heisenberg spin-$\frac{1}{2}$ finite chain, *Nucl.Phys.* **B554**, 647-678, (1999).

N.Kitanine, J.M.Maillet, and V.Terras, Correlation functions of the XXZ Heisenberg spin-$\frac{1}{2}$ chain in a magnetic field, *Nucl.Phys.* **B567**, 554-582, (2000).

J.M.Maillet and V.Terras, On the quantum inverse scattering problem, *Nucl.Phys.* **B575**, 627-644, (2000).

N.Kitanine, J.M.Maillet, N.A.Slavnov, and V.Terras, Spin-spin correlation functions of the $XXZ-\frac{1}{2}$ Heisenberg chain in a magnetic field, *Nucl.Phys.* **B641**, 487-518, (2002).

[5] B.Davies, O.Foda, M.Jimbo, T.Miwa , and A.Nakayashiki, Diagonalization of the XXZ Hamiltonian by vertex operator, *Commun.Math.Phys.***151**, 89-153, (1993).

M.Jimbo, K.Miki, T.Miwa, and A.Nakayashiki, Correlation functions of the XXZ model for $\Delta < -1$, *Phys.Lett.* **A168**, 256-263, (1992).

M.Jimbo and T.Miwa, *Algebraic analysis of solvable lattice model*, CBMS Regional Conference Series in Mathematics **85**, AMS, 1994.

[6] N.Reshetikhin, Calculation of the norm of Bethe vectors in models with $SU(3)$-symmetry, *J. Soviet Math.* **46**, 1694-1706, (1989).

M.Gaudin, *La fonction d'onde de Bethe*, Masson, Paris, 1983.

V.E.Korepin, Calculation of norms of Bethe wave functions, *Comm. Math. Phys.* **86**, 391-418, (1982).

[7] M.Wheeler, Multiple integral formulae for the scalar product of on-shell and off-shell Bethe vectors in $SU(3)$-invariant models *Nucl.Phys.* **B875**, 186-212, (2013).

S.Belliard, S.Pakuliak, E.Ragoucy, N.A.Slavnov, Bethe vectors of quantum integrable models with $GL(3)$ trigonometric R-matrix, Preprint, (2013), [arXiv:1304.7602].

[8] I.B.Frenkel and N.Jing, Vertex representations of quantum affine algebras, *Proc.Nat.Acad.Sci.USA***85** 9373-9377 (1988).

[9] Y.Koyama, Staggered polarization of vertex models with $U_q(\widehat{\mathrm{sl}(n)})$-symmetry, *Comm.Math.Phys.* **164**, 277-291, (1994).

[10] N.Jing, S.Kang, and Y.Koyama, Vertex operators of quantum affine Lie algebras $U_q(D_n^{(1)})$, *Comm.Math.Phys.* **174**, 367-392, (1995).

[11] D.Bernard, Vertex operator representations of the quantum affine algebra $U_q(B_r^{(1)})$, *Lett.Math.Phys.***17**, 239-245, (1989).

[12] K.Kimura, J.Shiraishi, and J.Uchiyama, A level-one representation of the quantum affine superalgebra $U_q(\widehat{sl}(M+1|N+1))$, *Comm.Math.Phys.* **188**, 367-378, (1997).

Y-Z.Zhang, Level-one representations and vertex operators of quantum affine superalgebra $U_q(\widehat{gl}(N|N))$, *J.Math.Phys.* **40**, 6110-6124, (1999).

[13] E.K.Sklyanin, Boundary conditions for integrable quantum systems, *J. Phys.* **A 21**, 2375-2389, (1988).

[14] R.A.Pimenta and A.Lima-Santos, Algebraic Bethe ansatz for the six vertex model with upper triangular K-matrices, *J.Phys.***A46** (2013) 455002, `arXiv:1308.4446`.

[15] R.Nepomechie, Bethe ansatz solution of the open XXZ chain with nondiagonal boundary terms, Special issue on recent advances in the theory of quantum integrable systems, *J. Phys.* **A37**, 433-440, (2004).

[16] J.Cao, H-Q. Lin, K-J. Shi, and Y.Wang, Exact solution of XXZ spin chain with unparallel boundary fields, *Nucl.Phys.* **B663**, 487-519, (2003).

W-L. Yang and Y-Z. Zhang, On the second reference state and complete eigenstates of the open XXZ chain, *JHEP* **04**, 044 (11 pp.), (2007).

[17] R.Murgan, R.I.Nepomechie, and C.Shi, Boundary energy of the open XXZ chain from new exact solutions, *Ann. Henri Poincaré* **7**, 1429-1448, (2006).

[18] N.Kitanine, K.K.Kozlowski, J.M.Maillet, G.Niccoli, N.A.Slavnov, and V.Terras, Correlation functions of the open XXZ chain. I., *J. Stat. Mech.* P10009 (37 pp.), (2007).

[19] N.Kitanine, K.K.Kozlowski, J.M.Maillet, G.Niccoli, N.A.Slavnov, and V.Terras, Correlation functions of the open XXZ chain. II., *J. Stat. Mech.*, P07010 (33 pp.) (2008).

[20] P.Baseilhac and K.Koizumi, A new (in)finite-dimensional algebra for quantum integrable models, *Nucl.Phys.***B720**, 325-347, (2005).

P.Baseilhac and K.Koizumi, A deformed analogue of Onsager's symmetry in the XXZ open spin chain, *J.Stat.Mech.* **0510** (2005) P005, `arXiv:hep-th/0507053`.

P.Baseilhac and K.Koizumi, Exact spectrum of the XXZ open spin chain from the q-Onsager algebra representation theory, *J. Stat. Mech.*, P09006 (27pp.), (2007).

P.Baseilhac, The q-deformed analogue of the Onsager algebra : beyond the Bethe ansatz approach, *Nucl.Phys.***B754**, 309-328, (2006).

[21] P.Baseilhac and S.Belliard, The half-infinite XXZ chain in Onsager's approach, *Nucl.Phys.***B873** 550-583 (2013).

[22] P.Baseilhac and K.Shigechi, A new current algebra and the reflection equation, *Lett.Math.Phys.***92**, 47-65, (2010).

P.Baseilhac and S.Belliard, Central extension of the reflection equations and an analog of Miki's formula, J.Phys.**A44**, 415205 (2011), arXiv:1104.1591.

[23] M.Jimbo, R.Kedem, T.Kojima, H.Konno, and T.Miwa, XXZ chain with a boundary, *Nucl.Phys.***B441** [FS] 437-470, (1995).

M.Jimbo, R.Kedem, H.Konno, T.Miwa, and R.Weston, Difference equations in spin chains with a boundary, *Nucl.Phys.* **B448**, 429-456, (1995).

[24] P.Baseilhac and T.Kojima, Correlation functions of the half-infinite XXZ spin chain with a triangular boundary, *Nucl.Phys.***B880**, 378-413, (2014), arXiv:1309.7785 .

[25] T.Kojima, The massless XXZ chain with a boundary, *Int.J.Mod.Phys.* **A16**, 409-424, (2001).

[26] M. Kasatani, Boundary quantum Knizhnik-Zamolodchikov equation, in: New trends in quantum integrable systems, 157-171, World Sci. Publ., Hackensack, NJ, (2011).

[27] N.Reshetikhin, J.Stokman and B.Vlaar, Reflection quantum Knizhnik-Zamolodchikov equations and Bethe vectors, arXiv:1305.1113;
J.Stokman and B.Vlaar, Koornwinder polynomials and the XXZ spin chain, arXiv:1310.5545.

[28] H.J.de Vega and A.Gonza'lez-Ruiz, Boundary K-matrices for the XYZ, XXZ and XXX spin chains, *J.Phys.* **A27**, 6129-6137, (1994).

[29] S.Ghoshal and Al.Zamolodchikov, Boundary S matrix and boundary state in two-dimensional integrable quantum field theory, *Int.J.Mod.Phys.* **A9**, 3841-3885, (1994).

[30] R.J.Baxter, Variational approximations for square lattice models in statistical mechanics, *J.Stat.Phys.* **19**, 461-478, (1978).

[31] M.Jimbo, T.Miwa, and A.Nakayashiki, Difference equations for the correlation functions of the eight-vertex model, *J. Phys.* **A26**, 2199-2209,(1993).

[32] O.Foda and T.Miwa, Corner transfer matrices and quantum affine algebras, *Int.J.Mod.Phys.* **A7** Suppl.1A, 279-302, (1992).

[33] H.Furutsu and T.Kojima, The $U_q(\widehat{\mathrm{sl}}_n)$ analogue of the XXZ chain with a boundary, *J.Math.Phys.* **41**, 4413-4436, (2000).

[34] W-L.Yang and Y-Z.Zhang, Izergin-Korepin model with a boundary, *Nucl.Phys.* **B596**, 495-512, (2001).

T.Kojima, A remark on ground state of boundary Izergin-Korepin model, *Int.J.Mod.Phys.* **A26**, 1973-1989, (2011).

[35] T.Kojima, Diagonalization of transfer matrix of supersymmetry $U_q(\widehat{sl}(M+1|N+1))$ chain with a boundary, *J.Math.Phys.***54**, 043507 (40 pp.), (2013).

[36] P.Baseilhac and S.Belliard, Generalized q-Onsager algebras and boundary affine Toda field theories, *Lett.Math.Phys.***93** (2010) 213-228, arXiv:0906.1215.

[37] J.Cao, W-L.Yang, K.Shi, Y.Wang, Off-diagonal Bethe ansatz solutions of the anisotropic spin-1/2 chains with arbitrary boundary fields, Nucl. Phys. **B 877** [FS] (2013) 152-175, arXiv:1307.2023.

[38] S.Belliard and N.Crampé, Heisenberg XXX model with general boundaries : Eigenvectors from Algebraic Bethe ansatz, SIGMA **9** (2013) 072, arXiv:1309.6165.

[39] W.Galleas, *Functional relations from the Yang-Baxter algebra: Eigenvalues of the XXZ model with non-diagonal twisted and open boundary conditions*, Nucl.Phys. **B790** (2008), 524-542, arXiv:1002.1623.

[40] G.Niccoli, Antiperiodic spin-$\frac{1}{2}$ XXZ quantum chains by separation of variables: complete spectrum and form factors, *Nucl.Phys.***B870**, 397-420, (2013).

G.Niccoli, Non-diagonal open spin-$\frac{1}{2}$-XXZ quantum chains by separation of variables: complete spectrum and matrix elements of some quasi-local operators, *J.Stat.Mech.*, P10025, (42pp.), (2012).

S.Faldella, N.Kitanine, and G.Niccoli, Complete spectrum and scalar products for the open spin-$\frac{1}{2}$ XXZ quantum chains with non-diagonal boundary terms, arXiv:1307.3960.

[41] E.M.Rains, Transformations of elliptic hypergeometric integrals, *Annals. Math.* **171**, 169-243, (2010).

[42] E.T.Whittaker and G.N.Watson, A course of modern analysis, Cambridge University Press, 1902.